Communicable Disease, Morbidity
Statistics, 1952 Report
N.C. State Board of Health

C61
N81
1952
pt.1, c.3

The Library
of the
University of North Carolina

Collection of North Caroliniana
This book was presented
by
The Board
C614.1
N87v
1952, pt.1, c.3

ANNUAL REPORT

OF

PUBLIC HEALTH

STATISTICS SECTION

PART I

Communicable Disease Morbidity Statistics

1952

NORTH CAROLINA STATE BOARD OF HEALTH

DIVISION OF EPIDEMIOLOGY

ANNUAL REPORT

OF

PUBLIC HEALTH

STATISTICS SECTION

PART I

Communicable Disease Morbidity Statistics

1952

NORTH CAROLINA STATE BOARD OF HEALTH

DIVISION OF EPIDEMIOLOGY

J. W. R. NORTON, M. D.

SECRETARY

AND STATE HEALTH OFFICER

614.1
N87v
1952, pt.1
c.2

TABLE OF CONTENTS

Table — Page

1. Estimated Population of North Carolina by Color and Sex, 1918-1952 — 9

2. Summary of All Reportable Communicable Diseases by year, 1918-1952 — 10

3. Reported Cases of Amebiasis (Place of Residence)
   - A. By county and color, 1951 and 1952, with month of report, 1952 — 13
   - B. By year of report and age, 1945-1952 — 15
   - C. By year of report, color, and sex with rates, 1945-1952 — 16

4. Reported Untreated Cases of Chancroid (Place of Report)
   - A. By county and color, 1951 and 1952, with month of report, 1952 — 17
   - B. By year of report and age, 1930-1952 — 20
   - C. By year of report, color, and sex with rates, 1930-1952 — 21

5. Reported Cases of Diphtheria (Place of Residence)
   - A. By county and color, 1951 and 1952, with month of onset, 1952 — 22
   - B. By year and age, 1930-1952 — 26
   - C. By year, color, and sex with rates, 1930-1952 — 27

6. Reported Cases of Encephalitis by County and Color, 1951 and 1952, with month of onset, 1952 (Place of Residence) — 28

7. Reported Cases of Gastro-Enteritis (Place of Residence)
   - A. By county, color, and type, 1952 onsets — 29
   - B. By year and age, 1952 — 32
   - C. By year, color, and sex with rates, 1952 — 32

8. Reported Untreated Cases of Gonorrhea (Place of Report)
   - A. By county and color, 1951 and 1952, with type and sex, 1952 reports — 33
   - B. By year of report and age, 1930-1952 — 38
   - C. By year of report, color, and sex with rates, 1930-1952 — 39

9. Reported Untreated Cases of Granuloma Inguinale (Place of Report)
   - A. By county and color, 1951 and 1952, with month of report, 1952 — 40
   - B. By year of report and age, 1945-1952 — 42
   - C. By year of report, color, and sex with rates, 1945-1952 — 43

10. Reported Cases of Infectious Jaundice (Place of Residence)
    - A. By county, color, and type, 1952 onsets — 44
    - B. By year and age, 1952 — 46
    - C. By year, color, and sex with rates, 1952 — 46

11. Reported Untreated Cases of Lymphogranuloma Venereum (Place of Report)
    - A. By county and color, 1951 and 1952, with month of report, 1952 — 47
    - B. By year of report and age, 1945-1952 — 49
    - C. By year of report, color, and sex with rates, 1945-1952 — 50

TABLE OF CONTENTS (Continued)

Table — Page

12. Reported Cases of Malaria (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 51
    B. By year and age, 1937-1952 52
    C. By year, color, and sex with rates, 1937-1952 53

13. Reported Cases of Measles (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 54
    B. By year and age, 1930-1952 58
    C. By year, color, and sex with rates, 1930-1952 59

14. Reported Cases of Meningitis (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 60
    B. By year and age, 1930-1952 64
    C. By year, color, and sex with rates, 1930-1952 65

15. Reported Cases of Paratyphoid Fever (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 66
    B. By year and age, 1930-1952 67
    C. By year, color, and sex with rates, 1930-1952 68

16. Reported Cases of Poliomyelitis (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 69
    B. By year and age, 1931-1952 73
    C. By year, color, and sex with rates, 1931-1952 74

17. Reported Cases of Rocky Mountain Spotted Fever (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 75
    B. By year and age, 1932-1952 77
    C. By year, color, and sex with rates, 1932-1952 78

18. Reported Cases of Scarlet Fever (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 79
    B. By year and age, 1930-1952 82
    C. By year, color, and sex with rates, 1930-1952 85

19. Reported Cases of Septic Sore Throat (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 86
    B. By year and age, 1930-1952 88
    C. By year, color, and sex with rates, 1930-1952 89

20. Reported Cases of Smallpox (Place of Residence)
    A. By year of report and age, 1930-1952 90
    B. By year, color, and sex with rates, 1930-1952 91

21. Reported Untreated Cases of Syphilis (Place of Report)
    A. By county and color, 1951 and 1952, with stage and sex, 1952 92
    B. By year of report and age, 1930-1952 96
    C. By year of report, color, and sex with rates, 1930-1952 97
    D. By color, stage, and sex with rates, 1940-1952 98
    E. Congenital syphilis, previously untreated, by color showing year of report, 1945-1952 and year of birth, 1925-1952: North Carolina, all report sources 102

TABLE OF CONTENTS (Continued)

Table — Page

22. Reported Cases of Tetanus (Place of Residence)
    A. By county and color, with month of onset, 1952 104
    B. By year and age, 1952 105
    C. By year, color, and sex with rates, 1952 105

23. Reported Cases of Trachoma (Place of Residence)
    A. By year of report and age, 1931-1952 106
    B. By year, color, and sex with rates, 1931-1952 107

24. Reported Cases of Tuberculosis, All Forms (Place of Residence)
    A. By county, color, and type, 1951 and 1952, with stage and activity, 1952 108
    B. By year of report and age, 1933-1952 114
    C. By year of report, color, and sex with rates, 1933-1952 115

25. Reported Cases of Tularemia (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 116
    B. By year and age, 1930-1952 118
    C. By year, color, and sex with rates, 1930-1952 119

26. Reported Cases of Typhoid Fever (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 120
    B. By year and age, 1930-1952 122
    C. By year, color, and sex with rates, 1930-1952 123

27. Reported Cases of Typhus Fever, Murine (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 124
    B. By year and age, 1930-1952 125
    C. By year, color, and sex with rates, 1930-1952 126

28. Reported Cases of Undulant Fever (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 127
    B. By year and age, 1932-1952 129
    C. By year, color, and sex with rates, 1932-1952 130

29. Reported Cases of Whooping Cough (Place of Residence)
    A. By county and color, 1951 and 1952, with month of onset, 1952 131
    B. By year and age, 1930-1952 135
    C. By year, color, and sex with rates, 1930-1952 136

INTRODUCTION

The twenty-first bulletin of North Carolina communicable disease morbidity statistics, covering the calendar year 1952, is published as Part I of the Annual Report of the Public Health Statistics Section. Part II, which contains natality and mortality data, will be issued later in the year.

Information regarding the diseases included in this report is shown by date of onset for all diseases except the five venereal diseases, tuberculosis, and amebiasis, which appear by date of report. Reports of all diseases having their onset in 1952 and received during the period January 1, 1952—January 31, 1953, plus unknown onsets received in 1952 are included except for those diseases tabulated by date of report. Data regarding the latter are confined to reports received during 1952, regardless of onset date.

All diseases included in this report are tabulated by place of residence, except the five venereal diseases which are shown by place of report. The venereal disease case reports are allocated to the county of report unless reported by a hospital or institution, in which case they are assigned to the county of residence.

Except for the first two tables, all information is grouped by disease. Table 2 presents a summary of reportable diseases in all previous years. As of 1952, five new reportable diseases were added. They were botulism, gastro-enteritis, infectious jaundice, Q fever, and tetanus. No cases were reported for botulism and Q fever. One disease, formerly shown as amebic dysentery, will hereafter be shown as amebiasis. Gastro-enteritis in Table 7-A, and infectious jaundice in Table 10-A are shown by type without month of onset. These data may be obtained upon request. Tetanus is reported for the first time and is shown in Table 22.

It may be of interest to note that poliomyelitis, with or without paralysis, was classified and tabulated as such for the first time in 1952. Although such details are not published in this report, data will be made available upon request.

All rates shown are per 100,000 population. Those instances in which counties are not shown in the various tables indicate that no cases were reported.

For the first time since 1946 malaria reports reached a new high of 305 resident cases. Of this number, 302 were traceable to military origin. In other words, all such cases were acquired by individuals while in service but were reported for the first time in 1952.

The diphtheria rate showed definite improvement during 1952 as compared to 1951. Only 202 cases were reported during 1952, a decline of 45 per cent under the number of cases reported (373) in 1951. The North Carolina diphtheria rate was 9.0 per 100,000 population in 1951 and was 4.8 per 100,000 in 1952.

Infectious jaundice, made a reportable disease in January 1952, occurred throughout the state in endemic form and was reported as highly prevalent in a few areas. A total of 675 cases was reported by card. Having been made a reportable disease for the first time during the year, and based on estimates submitted by local health officers in several counties, it is considered to be probable that many mild cases of this disease were not reported by attending physicians.

TABLE 1.—ESTIMATED POPULATION OF NORTH CAROLINA BY COLOR AND SEX, 1918-1952*

Year	All classes			White			Other		
	Total	Male	Female	Total	Male	Female	Total	Male	Female
1918	2,521,891	1,225,455	1,296,436	1,747,798	849,618	898,180	774,093	375,837	398,256
1919	2,534,828	1,255,353	1,279,475	1,746,614	871,112	875,502	788,214	384,241	403,973
1920	2,587,536	1,294,396	1,293,140	1,808,713	912,302	896,411	778,823	382,094	396,729
1921	2,650,711	1,328,295	1,322,416	1,859,539	940,314	919,225	791,172	387,981	403,191
1922	2,699,923	1,342,609	1,357,314	1,895,420	948,146	947,274	804,503	394,463	410,040
1923	2,761,364	1,373,738	1,387,626	1,942,055	972,207	969,848	819,309	401,531	417,778
1924	2,830,151	1,420,367	1,409,784	1,997,964	1,012,930	985,034	832,187	407,437	424,750
1925	2,894,514	1,444,843	1,449,671	2,031,248	1,022,417	1,008,831	863,266	422,426	440,840
1926	2,959,392	1,470,594	1,488,798	2,096,611	1,048,565	1,048,046	862,781	422,029	440,752
1927	3,027,006	1,495,951	1,531,055	2,147,539	1,067,478	1,080,061	879,467	428,473	450,994
1928	3,082,380	1,534,590	1,547,790	2,175,492	1,092,479	1,083,013	906,888	442,111	464,777
1929	3,133,082	1,558,156	1,574,926	2,208,369	1,107,772	1,100,597	924,713	450,384	474,329
1930	3,171,108	1,575,791	1,595,317	2,236,107	1,120,844	1,115,263	935,001	454,947	480,054
1931	3,206,641	1,593,707	1,612,934	2,266,838	1,136,266	1,130,572	939,803	457,441	482,362
1932	3,264,916	1,622,976	1,641,940	2,314,009	1,159,875	1,154,134	950,907	463,101	487,806
1933	3,318,377	1,649,920	1,668,457	2,357,442	1,181,608	1,175,834	960,935	468,312	492,623
1934	3,359,276	1,670,579	1,688,697	2,391,871	1,198,767	1,193,104	967,405	471,812	495,593
1935	3,380,227	1,681,169	1,699,058	2,409,930	1,207,605	1,202,325	970,297	473,564	496,733
1936	3,400,525	1,690,503	1,710,022	2,427,312	1,215,163	1,212,149	973,213	475,340	497,873
1937	3,432,099	1,705,708	1,726,391	2,453,834	1,227,533	1,226,301	978,265	478,175	500,090
1938	3,475,530	1,727,452	1,748,078	2,490,752	1,245,657	1,245,095	984,778	481,795	502,983
1939	3,531,938	1,755,416	1,776,522	2,536,869	1,268,107	1,268,762	995,069	487,309	507,760
1940	3,571,623	1,772,990	1,798,633	2,567,635	1,281,762	1,285,873	1,003,988	491,228	512,760
1941	3,670,017	1,820,328	1,849,689	2,635,072	1,313,866	1,321,206	1,034,945	506,462	528,483
1942	3,665,373	1,803,364	1,862,009	2,631,738	1,301,208	1,330,530	1,033,635	502,156	531,479
1943	3,683,037	1,789,956	1,893,081	2,637,054	1,281,697	1,355,357	1,045,983	508,259	537,724
1944	3,594,865	1,703,966	1,890,899	2,573,923	1,211,469	1,362,454	1,020,942	492,497	528,445
1945	3,514,297	1,637,662	1,876,635	2,512,722	1,156,203	1,356,519	1,001,575	481,459	520,116
1946	3,640,645	1,780,275	1,860,370	2,606,702	1,277,866	1,328,836	1,033,943	502,409	531,534
1947	3,718,000	1,825,538	1,892,462	2,662,088	1,312,454	1,349,634	1,055,912	513,084	542,828
1948	3,798,000	1,864,818	1,933,182	2,719,368	1,340,694	1,378,674	1,078,632	524,124	554,508
1949	3,864,000	1,897,224	1,966,776	2,766,624	1,363,946	1,402,678	1,097,376	533,278	564,098
1950	4,061,929	2,017,105	2,044,824	2,983,121	1,489,840	1,493,281	1,078,808	527,265	551,543
1951	4,123,218	2,047,620	2,075,598	3,035,057	1,515,850	1,519,207	1,088,161	531,770	556,391
1952	4,172,249	2,072,032	2,100,217	3,076,606	1,536,658	1,539,948	1,095,643	535,374	560,269

* Source—"Vital Statistics Rates in the United States 1900–1940" and "Current Population Reports," Bureau of the Census, 1941–1949. The census years 1920, 1930, 1940, 1950 by color and sex are not estimated; all other totals are estimated. The color and sex estimates for 1951, 1952 were obtained by extrapolation.

TABLE 2.—SUMMARY OF ALL REPORTABLE

Disease	1918	1919	1920	1921	1922	1923	1924	1925	1926	1927	1928	1929
Anthrax	—	—	—	—	—	—	—	—	—	—	—	—
Chancroid	+	429	463	222	260	210	169	170	191	151	106	72
Chicken pox	298	1,575	2,163	3,341	4,573	3,974	5,981	3,845	4,941	4,818	4,695	4,997
Cholera	—	—	—	—	—	—	—	—	—	—	—	—
Dengue	+	+	+	+	+	+	+	+	+	+	+	+
Diphtheria	1,366	3,519	3,422	5,136	8,136	4,671	4,095	3,437	3,198	3,034	3,826	4,337
Dysentery, bacillary	1	128	9	—	—	—	—	—	22	2	—	—
Encephalitis	+	+	+	+	+	+	+	+	+	+	+	+
German measles	+	43	75	72	130	159	134	106	4,481	389	172	1,267
Glanders	+	+	+	+	+	+	+	+	+	+	+	+
Gonorrhea**	+	4,893	5,361	3,475	4,061	3,023	2,561	2,272	2,113	2,016	1,528	2,130
Granuloma inguinale	+	+	+	+	+	+	+	+	+	+	+	+
Influenza	+	14,128	76,306	—	—	—	—	—	—	—	—	168
Leprosy	+	+	+	+	+	+	+	+	+	+	+	+
Lymphogranuloma venereum	+	+	+	+	+	+	+	+	+	+	+	+
Malaria	+	+	+	+	+	+	+	+	+	+	+	+
Measles	9,158	5,725	3,621	11,539	2,042	52,066	34,938	1,002	7,952	32,882	60,543	1,209
Meningococcus meningitis	151	105	96	70	54	46	30	26	24	26	21	88
Ophthalmia neonatorum	12	33	27	19	16	15	5	8	6	5	6	12
Paratyphoid fever	1	26	2	—	—	—	—	—	—	—	—	13
Plague	—	—	—	—	—	—	—	—	—	—	—	—
Poliomyelitis	23	42	25	36	30	16	16	79	112	24	51	133
Psittacosis	—	—	—	—	—	—	—	—	—	—	—	—
Rabies	—	—	—	—	—	—	—	—	—	—	—	2
Rocky Mountain spotted fever	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever	864	1,512	1,663	2,585	3,153	2,418	2,255	1,856	2,281	2,638	2,828	3,057
Septic sore throat	127	371	218	199	281	150	143	104	92	127	104	141
Smallpox	983	2,322	2,961	2,513	1,409	3,352	3,845	1,920	1,594	1,702	2,419	589
Syphilis	+	2,084	2,213	2,637	4,671	3,559	3,153	3,161	3,044	3,224	2,846	4,457
Trachoma	20	16	15	5	12	18	5	32	3	2	3	2
Tuberculosis, all forms	3,514	2,306	3,295	2,523	1,356	1,851	2,381	2,355	1,921	2,271	2,578	2,858
Tularemia	—	—	—	—	—	—	—	—	—	—	—	4
Typhoid fever	3,461	2,956	1,856	2,099	1,960	1,577	1,318	1,192	1,502	1,280	1,073	861
Typhus fever, murine	—	—	2	—	—	—	—	—	—	—	—	3
Typhus fever, European	—	—	—	—	—	—	—	—	—	—	—	—
Undulant fever	—	—	—	—	—	—	—	—	—	—	—	—
Vincent's infection	+	+	+	+	+	+	+	+	+	+	+	+
Whooping cough	11,645	5,669	8,465	9,518	6,905	16,075	12,945	4,296	11,099	19,996	5,002	12,434
Yellow fever	—	—	—	—	—	—	—	—	—	—	—	—

* 1918–1947 by year of report; 1948–1952 by year of onset, except the venereal diseases, tuberculosis, and amebiasis which are by year of report.

— Disease was reportable but none was reported.

\+ Disease not reportable.

** Gonorrhea totals include ophthalmia neonatorum.

† Data not obtainable.

Communicable Diseases by Year of Report, 1918-1952*

1930	1931	1932	1933	1934	1935	1936	1937	1938	1939	1940	1941	1942	1943	1944
—	1	—	—	2	—	—	—	—	—	—	1	—	—	—
114	94	88	35	52	73	110	148	102	321	264	223	635	663	451
6,982	4,856	4,226	4,167	4,540	3,843	3,179	5,409	5,316	3,765	4,360	4,726	5,063	4,698	3,782
—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
+	+	+	+	+	1	1	—	—	—	—	—	—	—	—
3,248	3,156	1,895	2,497	2,114	1,720	2,347	2,056	2,442	2,368	1,125	1,629	1,187	801	665
—	—	1	1	—	4	3	19	10	22	9	12	35	130	12
+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
967	9,384	394	187	604	589	3,326	2,855	228	244	168	12,769	331	2,374	1,316
+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
2,450	2,298	2,130	2,176	2,164	1,826	2,159	2,894	3,045	3,815	4,616	5,834	9,637	10,870	10,892
+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
856	4,739	3,841	7,215	1,564	3,215	2,830	1,979	498	1,544	2,113	5,837	673	1,404	1,294
+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
+	+	+	+	+	+	+	876	636	629	629	237	248	183	154
1,376	15,499	13,672	16,197	55,764	10,683	1,395	7,937	49,614	20,031	3,685	27,764	21,581	5,217	27,655
153	123	71	66	50	109	196	169	93	59	34	46	49	478	308
14	12	18	9	14	19	22	17	16	15	22	29	17	11	†
21	27	16	12	15	26	20	16	18	16	3	11	7	6	7
—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
96	104	54	30	45	675	52	106	50	116	74	168	82	36	861
—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
2	—	—	1	—	4	3	2	—	1	1	—	2	—	1
—	—	1	27	34	21	32	27	42	41	32	20	35	36	59
3,096	3,461	2,868	3,662	2,852	2,072	1,921	2,073	2,228	2,599	2,442	2,165	2,570	2,854	2,239
114	153	142	123	104	116	115	125	148	116	88	139	109	201	47
556	63	72	35	18	24	19	11	35	15	6	4	7	21	4
5,106	5,144	4,365	4,570	5,201	5,189	6,086	12,299	20,714	30,985	21,333	17,560	16,713	14,125	9,804
—	6	—	5	1	6	1	—	—	—	8	4	—	1	—
3,201	2,638	2,488	2,524	2,047	1,996	3,008	2,632	2,607	2,366	2,084	2,425	1,902	1,633	1,804
7	3	10	28	14	15	10	13	26	26	12	22	13	14	10
1,000	991	823	684	464	645	518	493	485	343	238	197	181	111	122
16	14	24	46	23	51	33	68	81	122	65	78	100	157	236
—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
—	—	4	15	19	31	23	26	29	25	11	12	9	7	14
+	+	+	+	+	+	+	+	29	14	134	85	29	66	174
10,093	6,992	12,137	7,155	14,299	10,075	1,566	7,782	14,908	9,321	5,500	10,932	5,557	8,408	6,355
—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

TABLE 2.—SUMMARY—*Continued*

Disease	1945	1946	1947	1948	1949	1950	1951	1952
Amebiasis	21	19	18	29	91	132	96	44
Anthrax	—	—	—	—	—	—	—	—
Botulism	+	+	+	+	+	+	+	—
Chancroid	447	513	531	381	332	265	253	256
Cholera	—	—	—	—	—	—	—	—
Dengue	—	1	—	—	—	—	—	—
Diphtheria	1,475	590	751	485	550	499	373	202
Encephalitis	5	3	5	6	4	9	9	4
Gastro-enteritis	+	+	+	+	+	+	+	219
Glanders	—	—	—	—	—	—	—	—
Gonorrhea**	13,162	16,082	14,169	14,962	16,173	14,930	15,779	15,895
Granuloma inguinale	75	101	99	98	66	82	58	38
Infectious jaundice	+	+	+	+	+	+	+	675
Leprosy	1	—	—	—	—	—	—	1
Lymphogranuloma venereum	231	153	120	176	129	91	81	62
Malaria	554	369	137	147	53	36	82	305
Measles	1,194	9,011	4,978	1,353	17,330	4,964	3,315	4,511
Meningococcus meningitis	179	120	95	73	87	116	151	197
Ophthalmia neonatorum	†	6	7	3	11	8	9	14
Paratyphoid fever	5	6	7	13	8	7	5	4
Plague	—	—	—	—	—	—	—	—
Poliomyelitis	159	157	300	2,516	229	756	314	538
Psittacosis	—	1	—	—	—	—	—	—
Q-fever	+	+	+	+	+	+	+	—
Rabies	1	—	2	—	—	—	1	—
Rocky Mountain spotted fever	57	66	88	73	79	70	61	38
Scarlet fever	3,248	1,530	1,167	1,094	1,793	1,885	2,236	2,222
Septic sore throat	101	51	44	27	45	41	29	33
Smallpox	1	2	2	3	—	—	—	—
Syphilis	8,125	9,233	8,724	7,313	6,699	5,146	4,080	3,709
Tetanus	+	+	+	+	+	+	+	26
Trachoma	1	—	—	—	—	—	—	—
Trichinosis	+	+	+	+	+	+	+	1
Tuberculosis, all forms	3,392	3,466	3,591	3,274	3,402	3,653	3,105	2,326
Tularemia	27	45	74	46	38	25	17	19
Typhoid fever	82	54	47	56	53	58	44	41
Typhus fever, murine	159	82	51	43	26	12	19	12
Typhus fever, European	—	—	—	—	—	—	—	—
Undulant fever	15	9	21	16	25	21	28	21
Whooping cough	5,834	3,391	2,983	1,932	1,398	3,352	1,961	471
Yellow fever	—	—	—	—	—	—	—	—

* 1918–1947 by year of report; 1948–1952 by year of onset, except the venereal diseases, tuberculosis, and amebiasis which are by year of report.

— Disease was reportable but none was reported.

\+ Disease not reportable.

** Gonorrhea totals include ophthalmia neonatorum.

† Data not obtainable.

Table 3A.—Reported Cases of Amebiasis by County and Color, 1951 and 1952, With Month of Report, 1952
(Place of Residence)

Area	1951 Reports				1952 Reports																
	Color				Color				Month of report												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	96	89	7	------	44	37	7	------	4	7	1	------	1	3	4	8	2	3	10	1	------
Bertie	1	------	1	------	1	------	1	------	------	------	------	------	------	------	------	------	------	1	------	------	------
Brunswick	2	2	------	------	5	4	1	------	------	1	------	------	------	1	2	------	------	1	------	------	------
Buncombe	------	------	------	------	1	1	------	------	------	------	------	------	------	------	------	1	------	------	------	------	------
Caldwell	------	------	------	------	1	------	1	------	------	------	------	------	------	------	------	------	------	1	------	------	------
Cleveland	------	------	------	------	1	1	------	------	------	1	------	------	------	------	------	------	------	------	------	------	------
Craven	2	2	------	------	1	1	------	------	------	------	------	------	------	------	------	1	------	------	------	------	------
Edgecombe	3	3	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Forsyth	1	------	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Granville	1	1	------	------	3	3	------	------	------	1	1	------	1	------	------	------	------	------	------	------	------
Halifax	3	3	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Haywood	9	9	------	------	1	1	------	------	1	------	------	------	------	------	------	------	------	------	------	------	------
Henderson	1	1	------	------	1	1	------	------	------	------	------	------	------	------	------	1	------	------	------	------	------
Jackson	35	34	1	------	6	6	------	------	1	------	------	------	------	------	------	4	1	------	------	------	------
Johnston	1	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Lee	1	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

TABLE 3A.—AMEBIASIS—*Continued*

Area	1951 Reports				1952 Reports																
	Color				Color				Month of report												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Lenoir	1		1		2	2				2											
Macon	14	14			3	3			1	1					1						
Madison	1	1																			
Martin	2	2																			
Mecklenburg					1		1										1				
Nash	3	3																			
New Hanover	3	3																			
Orange					1		1												1		
Pitt	3	3																			
Robeson	2		2		1		1													1	
Rowan					11	11								2					9		
Rutherford					1	1				1											
Sampson	1		1																		
Surry					1	1										1					
Swain	1	1																			
Transylvania	1	1																			
Wake	2	2			1	1									1						
Washington	2	2																			
Wilson					1		1		1												

Table 3B.—Reported Cases of Amebiasis by Year of Report and Age, 1945-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1945	21	------	------	------	------	------	------	------	------	------	------	------	1	1	7	4	3	2	2	------	------	1
1946	19	------	------	------	------	------	------	------	------	------	------	------	1	5	6	4	2	1	------	------	------	------
1947	18	------	1	------	------	------	------	------	------	------	------	------	------	5	5	2	1	3	------	------	------	1
1948	29	1	------	2	------	------	1	------	------	------	------	2	1	6	1	4	4	3	2	2	------	------
1949	91	2	------	2	------	1	------	------	1	------	------	7	2	8	10	11	29	7	5	1	------	5
1950	132	------	1	1	------	------	------	------	1	4	1	9	16	6	10	18	31	18	8	4	1	3
1951	96	------	------	------	------	------	------	1	1	2	1	4	4	5	11	15	21	16	11	3	1	------
1952	44	2	3	------	1	------	1	------	1	1	2	6	1	2	2	2	8	7	1	3	1	------

TABLE 3C.—REPORTED CASES OF AMEBIASIS BY YEAR OF REPORT, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1945-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1945	21	0.6	16	1.0	5	0.3	14	0.6	10	0.9	4	0.3	7	0.7	6	1.2	1	0.2	----------
1946	19	0.5	17	1.0	2	0.1	15	0.6	13	1.0	2	0.2	4	0.4	4	0.8	------	------	----------
1947	18	0.5	15	0.8	3	0.2	18	0.7	15	1.1	3	0.2	------	------	------	------	------	------	----------
1948	29	0.8	16	0.9	13	0.7	26	1.0	15	1.1	11	0.8	3	0.3	1	0.2	2	0.4	----------
1949	91	2.4	38	2.0	53	2.7	87	3.1	34	2.5	53	3.8	4	0.4	4	0.8	------	------	----------
1950	132	3.2	61	3.0	71	3.5	121	4.1	55	3.7	66	4.4	11	1.0	6	1.1	5	0.9	----------
1951	96	2.3	43	2.1	53	2.6	89	2.9	37	2.4	52	3.4	7	0.6	6	1.1	1	0.2	----------
1952	44	1.1	28	1.4	16	0.8	37	1.2	24	1.6	13	0.8	7	0.6	4	0.7	3	0.5	----------

Table 4A.—Reported Untreated Cases of Chancroid by County and Color, 1951 and 1952, With Month of Report, 1952
(Place of Report)

Area	1951 Reports				1952 Reports																
	Color				Color				Month of report												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	253	28	225		256	38	218		32	20	23	39	24	25	19	11	9	23	18	13	
Alamance	1		1		2		2						1	1							
Beaufort					1	1				1											
Bladen					1		1										1				
Buncombe	4	1	3		5	4	1			1						1		1	2		
Burke	2		2																		
Cabarrus	1	1																			
Carteret					1		1													1	
Cleveland					1		1						1								
Columbus	6		6		2		2		1											1	
Craven					8	6	2			2				1	1		1	2		1	
Cumberland	32	3	29		29	12	17		5	1	6	5	1	3	2	2		2	2		
Davidson	1		1																		
Davie	2		2																		
Duplin	3		3		1		1			1											
Durham	32	1	31		21	1	20		1	2	1	5	1		5	2	2	2			

TABLE 4A.—CHANCROID—*Continued*

Area	1951 Reports				1952 Reports															
	Color				Color				Month of report											
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Edgecombe	1		1		2		2				1									1	
Forsyth	13		13		11		11		2			1	2	1			1		4		
Franklin	2		2																		
Gaston	7		7																		
Granville					2		2		2												
Greene	1		1																		
Guilford	4		4		4		4		1			1		1				1			
Harnett	3		3		1		1		1												
Hoke	1		1																		
Johnston	3		3																		
Lee	1		1		1		1				1										
Lenoir	3	3			1		1											1			
McDowell	1		1																		
Mecklenburg	66	5	61		101	1	100		15	7	7	20	13	10	9	5	1	8	2	4	
Moore	2		2																		
Nash	1		1		2		2												1	1	
New Hanover	8		8		12	1	11		2	2	1			2	1		1	1	1	1	
Northampton	2		2																		
Onslow	19	13	6		13	9	4		2	2	2	3	2	2							
Perquimans					1		1												1		

Person	1		1																		
Pitt	3		3																		
Richmond	1		1		2		2			1			1								
Robeson					7		7				1	3		2						1	
Rowan	1		1		3		3				1								2		
Sampson	1		1		1		1											1			
Scotland	1	1			1		1										1				
Transylvania					1	1													1		
Vance	1		1		1		1									1					
Wake	8		8		2		2						1				1				
Wayne	5		5		10	2	8				1		1	1	1			3	1	2	
Wilson	9		9		5		5				1	1		1				1	1		

TABLE 4B.—REPORTED UNTREATED CASES OF CHANCROID BY YEAR OF REPORT AND AGE, 1930-1952 (Place of Report)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	114												14	34	18	11	6	4	2			25
1931	94											2	19	24	16	7	4	2	2			18
1932	88											1	19	28	14	3	10	2	2			9
1933	35												4	10	8	2	4	1				6
1934	52												9	16	8	4	5			1		9
1935	73					1	5						10	25	11	10	6	3				2
1936	110												22	41	22	10	9	4		1		1
1937	148											1	27	45	32	17	11	6	1			8
1938	102												23	32	12	9	14	2			1	9
1939	321											2	79	130	46	30	26	2	1			5
1940	264											2	65	107	38	24	21	2	1			4
1941	223							3	2	3	6	17	48	65	45	18	11	5				
1942	635											5	124	226	135	74	52	11	4	2		2
1943	663					1						3	154	272	128	62	35	7	1			
1944	451							1				6	120	170	84	34	26	9	1			
1945	447											5	119	148	87	49	27	6	2	3	1	
1946	513											4	100	237	106	36	20	6	1		2	1
1947	531											4	133	183	104	51	39	8	7	1		1
1948	381												86	148	82	28	29	7	1			
1949	332											3	74	118	56	34	30	13	1	1		2
1950	265											1	69	95	42	26	19	10	2	1		
1951	253											5	55	92	45	19	24	8	3	2		
1952	256											2	66	85	52	19	26	5	1			

TABLE 4C.—REPORTED UNTREATED CASES OF CHANCROID BY YEAR OF REPORT, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Report)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	114	3.6	104	6.6	10	0.6	60	2.7	56	5.0	4	0.4	46	4.9	40	8.8	6	1.2	8
1931	94	2.9	83	5.2	11	0.7	40	1.8	39	3.4	1	0.1	54	5.7	44	9.6	10	2.1	----------
1932	88	2.7	82	5.1	6	0.4	33	1.4	33	2.8	-------	-------	55	5.8	49	10.6	6	1.2	----------
1933	35	1.1	34	2.1	1	0.1	12	0.5	12	1.0	-------	-------	21	2.2	20	4.3	1	0.2	2
1934	52	1.5	50	3.0	2	0.1	22	0.9	22	1.8	-------	-------	28	2.9	26	5.5	2	0.4	2
1935	73	2.2	61	3.6	12	0.7	23	1.0	21	1.7	2	0.2	50	5.2	40	8.4	10	2.0	----------
1936	110	3.2	105	6.2	5	0.3	34	1.4	33	2.7	1	0.1	76	7.8	72	15.1	4	0.8	----------
1937	148	4.3	143	8.4	5	0.3	45	1.8	44	3.6	1	0.1	103	10.5	99	20.7	4	0.8	----------
1938	102	2.9	99	5.7	3	0.2	24	1.0	24	1.9	-------	-------	71	7.2	68	14.1	3	0.6	7
1939	321	9.1	278	15.8	43	2.4	51	2.0	43	3.4	8	0.6	270	27.1	235	48.2	35	6.9	----------
1940	264	7.4	241	13.6	23	1.3	40	1.6	39	3.0	1	0.1	224	22.3	202	41.1	22	4.3	----------
1941	223	6.1	175	9.6	48	2.6	27	1.0	22	1.7	5	0.4	196	18.9	153	30.2	43	8.1	----------
1942	635	17.3	538	29.8	97	5.2	58	2.2	47	3.6	11	0.8	577	55.8	491	97.8	86	16.2	----------
1943	663	18.0	590	33.0	73	3.9	48	1.8	43	3.4	5	0.4	615	58.8	547	107.6	68	12.6	----------
1944	451	12.5	349	20.5	102	5.4	76	3.0	45	3.7	31	2.3	375	36.7	304	61.7	71	13.4	----------
1945	447	12.7	335	20.5	112	6.0	99	3.9	82	7.1	17	1.3	348	34.7	253	52.5	95	18.3	----------
1946	513	14.1	426	23.9	87	4.7	86	3.3	76	5.9	10	0.8	427	41.3	350	69.7	77	14.5	----------
1947	531	14.3	429	23.5	102	5.4	93	3.5	85	6.5	8	0.6	438	41.5	344	67.0	94	17.3	----------
1948	381	10.0	310	16.6	71	3.7	45	1.7	40	3.0	5	0.4	336	31.2	270	51.5	66	11.9	----------
1949	332	8.6	246	13.0	86	4.4	39	1.4	32	2.3	7	0.5	293	26.7	214	40.1	79	14.0	----------
1950	265	6.5	214	10.6	51	2.5	37	1.2	35	2.3	2	0.1	228	21.1	179	33.9	49	8.9	----------
1951	253	6.1	202	9.9	51	2.5	28	0.9	27	1.8	1	0.1	225	20.7	175	32.9	50	9.0	----------
1952	256	6.1	200	9.7	56	2.7	38	1.2	37	2.4	1	0.1	218	19.9	163	30.4	55	9.8	----------

TABLE 5A.—REPORTED CASES OF DIPHTHERIA BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets															
	Color				Color				Month of onset											
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	373	189	184		202	112	89	1	17	13	13	14	9	8	6	16	42	30	20	14	
Alamance					8	5	3		1			2		3		2					
Alexander	2	2																			
Alleghany					1	1					1										
Anson	1		1		1		1									1					
Ashe	2	2																			
Beaufort	9	2	7		2	1	1		2												
Bertie	4	2	2		2	2											1	1			
Bladen	5	5																			
Brunswick	6	3	3																		
Buncombe	4	3	1		3	3			1						2						
Burke	7	7			1	1							1								
Cabarrus	4	4			3	3						2		1							
Caldwell	2	1	1		1	1				1											
Carteret	1		1																		
Caswell	11	1	10		1	1					1										

Catawba	1	1			1	1					1										
Cherokee	2	2																			
Clay	1	1																			
Cleveland	7	2	5		2		2				1					1					
Columbus	1	1																			
Craven	7	6	1		3	1	2			2						1					
Cumberland	2	2			2	1	1						1						1		
Currituck	1		1																		
Davidson	1	1			2	2									1	1					
Duplin	3	1	2		7	3	4		1		2						2	1		1	
Durham	4	4			3	2	1		1			1				1					
Edgecombe	8	3	5		7	2	5						1				4		2		
Forsyth	1	1			1	1													1		
Gaston	5	5			5	5			2	1								1		1	
Granville	3		3																		
Guilford	1	1																			
Halifax	24	4	20		14	4	9	1	1					2	1	1	3	6			
Harnett	2	2			1		1											1			
Haywood	11	10	1		1	1													1		
Henderson					5	5												4		1	
Hertford	15	7	8		1	1											1				
Hoke	3	2	1		3		3										2		1		
Iredell	8	6	2		2	2							1					1			
Jackson	2	2																			
Johnston	11	8	3		5	3	2									2	2		1		
Jones	4	4			4	4								1			1	1	1		
Lee	3	2	1		1		1											1			
Lenoir	17	7	10		12	9	3		1			1					3	2	5		
McDowell	1	1																			
Macon	6	6																			

TABLE 5A.—DIPHTHERIA—*Continued*

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Madison					7	7						2	3				1		1		
Martin	6	3	3		8	6	2				3		2	1						2	
Mecklenburg	2	2			7	4	3		1	1							1	2		2	
Mitchell					1	1				1											
Montgomery					1	1				1											
Moore	2	2			2	2				1							1				
Nash	4	2	2		4	1	3									1	3				
New Hanover	5	3	2		1		1										1				
Northampton	66	5	61		14	1	13		1	2	1					1	6			3	
Onslow	2	2																			
Orange	1		1		1	1													1		
Pasquotank	1		1		2		2									2					
Pender	2	2																			
Person	1	1			1		1									1					
Pitt	6	2	4		4		4										3		1		
Richmond	3	3																			
Robeson	4		4		10	2	8			1	2				2			1	3	1	
Rockingham					2	2			1									1			
Rowan	4	3	1		1	1			1												
Rutherford	2	2			3	2	1			1		1						1			

Sampson	2	2			4		4										1	3			
Scotland	8	4	4		6	2	4										2	3		1	
Transylvania	1	1																			
Union	1	1																			
Vance					1	1											1				
Wake	6	4	2		4	1	3			1						1				2	
Warren	2	1	1																		
Washington	22	15	7		4	4						4									
Watauga	1	1			3	3			1		1	1									
Wayne	3	1	2		2	1	1		2												
Wilkes					2	2											2				
Wilson	3	3			2	2											1		1		

TABLE 5B.—REPORTED CASES OF DIPHTHERIA BY YEAR* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	3,248	80	245	435	389	366	248	285	184	161	94	271	125	96	92	43	55	20	6	5	------	48
1931	3,156	80	205	341	386	359	297	250	212	157	102	293	120	93	72	51	61	22	12	3	------	40
1932	1,895	49	146	213	247	204	170	171	128	83	62	148	82	58	40	25	31	11	5	------	------	22
1933	2,497	58	151	269	274	285	214	239	160	117	104	233	93	100	47	52	52	8	5	1	------	35
1934	2,114	50	147	227	252	229	198	197	136	107	65	177	80	71	51	46	47	14	8	------	------	12
1935	1,720	52	128	156	186	197	159	178	107	88	67	148	63	61	44	27	33	4	3	------	------	19
1936	2,347	75	180	264	281	301	184	224	145	96	90	209	83	61	50	28	32	9	2	------	1	32
1937	2,056	53	165	223	261	262	190	175	151	105	67	156	64	51	41	28	24	12	6	------	1	21
1938	2,442	56	162	253	273	299	241	249	172	125	77	215	92	67	47	33	32	16	6	1	------	26
1939	2,368	59	176	251	293	305	235	223	150	130	78	207	69	45	38	26	30	12	4	2	------	35
1940	1,125	34	84	142	125	115	110	92	67	57	27	101	48	33	23	17	15	10	5	------	1	19
1941	1,629	33	71	183	175	192	176	162	123	99	68	153	57	29	33	14	27	9	2	------	1	22
1942	1,187	31	40	124	134	129	117	114	97	66	67	134	35	22	23	17	19	3	------	1	1	13
1943	801	26	31	83	116	100	72	61	46	43	35	90	25	22	14	12	10	3	------	------	------	12
1944	665	19	28	91	80	81	55	49	45	33	37	61	32	18	12	5	4	4	------	------	1	10
1945	1,475	26	34	150	184	192	177	119	89	75	51	192	57	32	34	25	21	5	1	------	1	10
1946	590	11	29	82	77	78	44	44	26	23	20	64	27	17	18	11	9	4	------	------	------	6
1947	751	30	39	85	72	82	72	51	53	42	28	96	29	25	22	7	8	3	4	2	------	1
1948	506	17	53	48	51	55	50	33	29	19	19	53	28	13	10	6	12	7	1	1	------	1
1949	550	19	59	47	56	60	51	34	25	33	17	72	20	23	11	9	10	1	------	3	------	--------
1950	499	15	33	50	66	51	37	41	25	20	24	71	21	13	9	10	9	3	------	1	------	--------
1951	373	4	26	30	43	44	38	15	15	14	22	56	24	15	7	7	7	4	1	1	------	--------
1952	202	5	16	17	26	25	20	8	6	9	4	28	10	10	2	4	8	3	1	------	------	--------

* 1930-1948 by year of report; 1949–1952 by year of onset.

TABLE 5C.—REPORTED CASES OF DIPHTHERIA BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	3,248	102.4	1,524	96.7	1,698	106.4	2,727	122.0	1,289	115.0	1,415	126.9	488	52.2	221	48.6	265	55.2	33
1931	3,156	98.4	1,436	90.1	1,718	106.5	2,758	121.7	1,253	110.3	1,505	133.1	379	40.3	175	38.3	204	42.3	19
1932	1,895	58.0	900	55.5	989	60.2	1,698	73.4	811	69.9	887	76.9	179	18.8	85	18.4	94	19.3	18
1933	2,497	75.2	1,237	75.0	1,258	75.4	2,227	94.5	1,124	95.1	1,101	93.6	264	27.5	109	23.3	155	31.5	6
1934	2,114	62.9	969	58.0	1,145	67.8	1,875	78.4	867	72.3	1,008	84.5	231	23.9	96	20.3	135	27.2	8
1935	1,720	50.9	805	47.9	915	53.9	1,525	63.3	719	59.5	806	67.0	191	19.7	86	18.2	105	21.1	4
1936	2,347	69.0	1,124	66.5	1,223	71.5	2,098	86.4	1,008	83.0	1,090	89.9	235	24.1	106	22.3	129	25.9	14
1937	2,056	59.9	944	55.3	1,112	64.4	1,753	71.4	816	66.5	937	76.4	293	30.0	127	26.6	166	33.2	18
1938	2,442	70.3	1,198	69.4	1,244	71.2	2,069	83.1	1,015	81.5	1,054	84.7	365	37.1	178	36.9	187	37.2	8
1939	2,368	67.0	1,073	61.1	1,242	69.9	1,999	78.8	933	73.6	1,066	84.0	316	31.8	140	28.7	176	34.7	53
1940	1,125	31.5	544	30.7	581	32.3	920	35.8	456	35.6	464	36.1	205	20.4	88	17.9	117	22.8	----------
1941	1,629	44.4	781	42.9	808	43 7	1,352	51.3	660	50.2	692	52.4	237	22.9	121	23.9	116	21.9	40
1942	1,187	32.4	543	30.1	615	33.0	995	37.8	468	36.0	527	39.6	163	15.8	75	14.9	88	16.6	29
1943	801	21.7	379	21.2	416	22.0	663	25.1	314	24.5	349	25.7	132	12.6	65	12.8	67	12.5	6
1944	665	18.5	317	18.6	331	17.5	510	19.8	243	20.1	267	19.6	138	13.5	74	15.0	64	12.1	17
1945	1,475	42.0	663	40.5	801	42.7	1,212	48.2	553	47.8	659	48.6	252	25.2	110	22.8	142	27.3	11
1946	590	16.2	285	16.0	299	16.1	458	17.6	231	18.1	227	17.1	126	12.2	54	10.7	72	13.5	6
1947	751	20.2	356	19.5	395	20.9	494	18.6	237	18.1	257	19.0	257	24.3	119	23.2	138	25.4	----------
1948	506	13.3	232	12.4	274	14.2	351	12.9	159	11.9	192	13.9	154	14.3	73	13.9	81	14.6	1
1949	550	14.2	268	14.1	282	14.3	376	13.6	192	14.1	184	13.1	172	15.7	74	13.9	98	17.4	2
1950	499	12.3	234	11.6	265	13.0	325	10.9	154	10.3	171	11.5	174	16.1	80	15.2	94	17.0	----------
1951	373	9.0	180	8.8	193	9.3	189	6.2	95	6.3	94	6.2	184	16.9	85	16.0	99	17.8	----------
1952	202	4.8	96	4.6	106	5.0	112	3.6	55	3.6	57	3.7	89	8.1	40	7.5	49	8.7	1

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 6.—REPORTED CASES OF ENCEPHALITIS BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	9	6	3		4	2	2						1			1	1		1		
Anson					1	1													1		
Chowan					1		1										1				
Cleveland	1	1																			
Cumberland	1	1																			
Gates	1		1																		
Onslow	3	1	2																		
Pitt	1	1																			
Polk	1	1																			
Transylvania					1	1										1					
Wake	1	1																			
Wilson					1		1						1								

Table 7A.—Reported Cases of Gastro-enteritis by County, Color, and Type, 1952 Onsets
(Place of Residence)

Area	1952 Onsets												
	Total	White	Other	Not stated	Salmonellosis			Shigellosis		Staphylococcus food poisoning		Other and unspecified	
					White	Other	Not stated	White	Other	White	Other	White	Other
North Carolina	219	163	55	1	27	19	1	23	5	21	2	92	29
Alamance	2		2			2							
Anson	2	1	1							1			1
Avery	1	1						1					
Bladen	5	3	2		1	1						2	1
Brunswick	17	11	6		1							10	6
Buncombe	3	3			1			1				1	
Craven	2	2			2								
Cumberland	9	7	2		2	2		3				2	
Duplin	1	1										1	
Durham	2	2			1							1	
Edgecombe	4	1	3		1								3
Forsyth	3	2	1		1							1	1
Gates	2	1	1					1					1
Greene	1	1								1			
Guilford	3	3			2					1			

TABLE 7A.—GASTRO-ENTERITIS—*Continued*

Area	1952 Onsets												
	Total	White	Other	Not stated	Salmonellosis			Shigellosis		Staphylococcus food poisoning		Other and unspecified	
					White	Other	Not stated	White	Other	White	Other	White	Other
Halifax	6	2	4		1	3				1			1
Harnett	2	2						1				1	
Henderson	1	1						1					
Hertford	2	2			2								
Iredell	2	2						2					
Johnston	1	1						1					
Lee	3	2	1		1	1		1					
Lenoir	4	1	3			2		1					1
Martin	1		1										1
Mecklenburg	7	2	5		1				4		1	1	
Mitchell	1	1										1	
Moore	3	2	1		2	1							
New Hanover	1		1										1
Northampton	2	2			2								
Onslow	14	14										14	
Pamlico	1	1			1								
Pender	1		1			1							
Person	1		1			1							
Pitt	2		2			2							
Polk	1	1								1			

Randolph	2	1	1					1					1
Richmond	1		1										1
Robeson	6	1	5						1			1	4
Rowan	3	1	2			2						1	
Stanly	73	69	4							16	1	53	3
Union	1	1										1	
Wake	14	12	1	1	3	1	1	9					
Watauga	2	2			2								
Wayne	1	1										1	
Wilson	3		3										3

TABLE 7B.—REPORTED CASES OF GASTRO-ENTERITIS BY YEAR AND AGE, 1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1952	219	47	8	12	4	3	5	4	2	1	1	10	15	18	16	14	19	23	10	4	1	2

TABLE 7C.—REPORTED CASES OF GASTRO-ENTERITIS BY YEAR, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1952 (Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1952	219	5.2	115	5.6	104	5.0	163	5.3	84	5.5	79	5.1	55	5.0	30	5.6	25	4.5	1

Table 8A.—Reported Untreated Cases of Gonorrhea by County and Color, 1951 and 1952, With Type and Sex, 1952 Reports (Place of Report)

Area	1951 Reports			1952 Reports			Site of infection															
							Genito-urinary				Ophthalmia								Other			
											Neonatorum				Other eye							
	Total	White	Other	Total	White	Other	White		Other		White		Other		White		Other		White		Other	
							M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F
North Carolina	15,779	1,977	13,802	15,895	2,037	13,858	1,393	638	8,535	5,258			8	6			3	2	2	4	21	25
Alamance	82	8	74	70	2	68	1	1	38	29			1									
Alexander	8	7	1	4	3	1		3		1												
Alleghany				2		2				1								1				
Anson	58		58	19	1	18	1		11	7												
Ashe	3	3		3	3			3														
Avery	1	1		1	1		1															
Beaufort	46	3	43	26		26			17	8				1								
Bertie	25		25	7		7			7													
Bladen	66		66	51		51			30	21												
Brunswick	25	2	23	23	6	17	5	1	15	2												
Buncombe	232	80	152	210	79	131	54	25	87	44												
Burke	54	30	24	54	36	18	25	11	14	4												
Cabarrus	251	55	196	284	88	196	50	38	147	49												
Caldwell	47	34	13	34	21	13	16	5	11	2												
Camden	34		34																			

TABLE 8A.—GONORRHEA—*Continued*

Area	1951 Reports			1952 Reports																				
							Site of infection																	
							Genito-urinary				Ophthalmia								Other			
											Neonatorum				Other eye							
							White		Other		White		Other		White		Other		White		Other	
	Total	White	Other	Total	White	Other	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F
Carteret	31	2	29	14		14			13	1												
Caswell	6		6	11		11			10	1												*
Catawba	97	35	62	94	31	63	21	9	55	7									1			1
Chatham	1		1	6		6			5	1												
Cherokee	2	2		1		1			1													
Chowan	4		4	15		15				15												
Cleveland	200	35	165	166	15	151	12	3	114	37												
Columbus	135	3	132	203	1	202		1	122	75			5									
Craven	556	128	428	632	95	537	88	7	402	133				2								
Cumberland	1,503	301	1,202	1,637	478	1,159	358	120	836	323												
Currituck	28	1	27	28		28			16	6											5	1
Dare				10		10			7	3												
Davidson	3	1	2																			
Davie	15		15	4	2	2	1	1	1	1												

Duplin	29	2	27	50	2	48		2	38	9											1	
Durham	1,065	44	1,021	1,093	30	1,063	19	10	648	411										1	2	2
Edgecombe	270	5	265	283	7	276	6	1	167	108							1					
Forsyth	1,463	81	1,382	1,497	111	1,386	57	54	673	712								1				
Franklin	29		29	48		48			31	17												
Gaston	223	92	131	172	73	99	52	20	77	22									1			
Gates	3		3	2		2			1	1												
Graham	3	3																				
Granville	85	3	82	101		101			47	54												
Greene	17		17	17		17			13	4												
Guilford	762	76	686	672	61	611	28	33	391	218											2	
Halifax	22	1	21	29	3	26	2		10	3										1		13
Harnett	121	4	117	166	6	160	3	3	118	42												
Haywood	7	7		11	9	2	2	7	1	1												
Henderson	20	2	18	29	6	23	4	2	21	2												
Hertford				1		1			1													
Hoke	78	1	77	61	5	56	4	1	42	14												
Hyde				3		3			2	1												
Iredell	63	2	61	44	6	38	5	1	21	17												
Jackson	2	2		3	2	1	2		1													
Johnston	45	2	43	35	9	26	5	4	14	12												
Jones	23	1	22	20		20			14	6												
Lee	3		3																			
Lenoir	151	4	147	230	16	214	5	11	76	135							1					2
Lincoln	14	7	7	4		4			2	2												
McDowell	18	4	14	27	19	8	15	4	6	2												
Macon	6	6		4	4		1	3														
Madison	1	1																				
Martin	33	2	31	32	2	30	1	1	24	6												
Mecklenburg	2,483	291	2,192	2,161	217	1,944	120	97	985	954			1	3							1	

TABLE 8A.—GONORRHEA—*Continued*

Area	1951 Reports			1952 Reports																			
							Site of infection																
							Genito-urinary				Ophthalmia								Other				
											Neonatorum				Other eye								
							White		Other		White		Other		White		Other		White		Other		
	Total	White	Other	Total	White	Other	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	
Mitchell	4	4	------	2	2	------	1	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	
Montgomery	12	2	10	11	1	10	------	1	5	5	------	------	------	------	------	------	------	------	------	------	------	------	
Moore	44	9	35	28	3	25	2	1	19	6	------	------	------	------	------	------	------	------	------	------	------	------	
Nash	359	5	354	290	2	288	2	------	175	113	------	------	------	------	------	------	------	------	------	------	------	------	
New Hanover	1,360	88	1,272	1,262	58	1,204	19	39	658	546	------	------	------	------	------	------	------	------	------	------	------	------	
Northampton	46	------	46	45	------	45	------	------	21	24	------	------	------	------	------	------	------	------	------	------	------	------	
Onslow	681	250	431	1,140	280	860	272	8	784	75	------	------	------	------	------	------	------	------	------	------	1	------	
Orange	109	1	108	72	2	70	1	1	41	28	------	------	------	------	------	------	1	------	------	------	------	------	
Pamlico	4	------	4	1	------	1	------	------	------	1	------	------	------	------	------	------	------	------	------	------	------	------	
Pasquotank	126	3	123	265	5	260	2	3	149	111	------	------	------	------	------	------	------	------	------	------	------	------	
Pender	22	1	21	39	------	39	------	------	21	18	------	------	------	------	------	------	------	------	------	------	------	------	
Perquimans	23	------	23	30	------	30	------	------	13	17	------	------	------	------	------	------	------	------	------	------	------	------	
Person	193	4	189	79	------	79	------	------	41	38	------	------	------	------	------	------	------	------	------	------	------	------	
Pitt	82	1	81	118	7	111	3	4	49	62	------	------	------	------	------	------	------	------	------	------	------	------	
Polk	5	2	3	1	------	1	------	------	1	------	------	------	------	------	------	------	------	------	------	------	------	------	

Randolph	42	8	34	42	10	32	8	2	21	11												
Richmond	63	19	44	37	8	29	4	4	19	10												
Robeson	136	15	121	118	12	106	9	2	66	40										1		
Rockingham	34	6	28	9	2	7		2	2	5												
Rowan	63	16	47	28	6	22	3	3	17	5												
Rutherford	34	12	22	24	1	23	1		15	8												
Sampson	17		17	20	1	19		1	4	3											6	6
Scotland	271	28	243	218	42	176	31	11	110	65			1									
Stanly	128	9	119	83	7	76	4	3	51	25												
Stokes				1		1				1												
Surry	96	59	37	120	73	47	35	38	24	23												
Swain	5		5	1		1				1												
Transylvania	1	1		3		3			2												1	
Tyrrell	4		4	6		6			1	5												
Union	21	5	16	8	3	5	2	1	2	3												
Vance	267	6	261	350	8	342	6	1	317	23										1	2	
Wake	461	20	441	419	19	400	12	7	232	168												
Warren	90		90	56		56			51	5												
Washington	3		3	2		2			2													
Watauga	1	1																				
Wayne	191	15	176	171	19	152	5	14	87	65												
Wilkes	3	3		9	8	1	5	3		1												
Wilson	218	7	211	375	3	372		3	151	221												
Yadkin				6	3	3		3	1	2												
Yancey	8	3	5	2	2		2															

TABLE 8B.—REPORTED UNTREATED CASES OF GONORRHEA BY YEAR OF REPORT AND AGE, 1930-1952 (Place of Report)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	2,450	17	3	5	10	11	5	10	7	10	2	21	464	819	459	238	202	57	28	2	2	78
1931	2,298	15	1	6	9	8	6	8	8	9	8	21	483	810	369	235	177	43	15	3	------	64
1932	2,130	19	------	4	4	6	7	4	11	12	4	20	454	730	379	170	193	55	13	2	1	42
1933	2,176	14	4	9	6	9	7	7	6	6	4	34	446	780	353	167	175	59	40	1	------	49
1934	2,164	13	1	7	12	10	6	8	7	11	5	31	385	565	586	189	168	40	19	3	------	98
1935	1,826	16	2	4	7	7	10	7	9	7	8	28	357	627	293	147	171	57	12	------	5	52
1936	2,159	16	3	8	8	9	29	43	28	36	25	72	415	684	334	192	164	41	15	4	------	33
1937	2,894	17	2	9	15	16	14	22	28	16	11	65	589	897	472	247	226	78	24	9	------	137
1938	3,045	17	4	18	14	19	11	21	19	17	17	62	681	980	510	226	222	68	27	7	1	104
1939	3,815	13	3	19	12	10	22	21	19	17	26	74	918	1,265	632	321	277	70	21	10	------	65
1940	4,616	21	5	23	15	12	26	25	23	21	31	90	1,110	1,526	765	390	325	85	25	12	------	86
1941	5,834	31	3	22	10	13	8	22	23	15	18	85	1,459	2,168	1,037	464	315	82	30	7	2	20
1942	9,637	24	------	7	3	11	8	11	10	13	9	94	1,971	4,140	1,757	813	597	85	25	7	2	50
1943	10,870	1	6	9	5	10	2	9	5	7	10	127	3,053	4,240	1,844	835	510	93	18	5	1	80
1944	10,892	10	5	6	5	12	10	5	9	8	2	102	2,849	4,393	2,089	856	437	68	17	5	4	------
1945	13,162	18	19	13	16	11	4	15	14	7	5	174	3,506	4,633	2,765	1,068	684	177	22	11	------	------
1946	16,082	------	12	16	11	11	7	12	7	9	11	159	4,130	6,826	2,969	1,036	643	143	31	10	2	37
1947	14,169	9	3	7	16	13	9	11	15	15	9	176	3,751	5,900	2,540	898	613	118	27	9	------	30
1948	14,962	6	8	11	12	12	7	13	13	16	20	167	3,799	6,263	2,761	965	694	133	35	12	2	13
1949	16,173	10	7	12	11	14	9	11	15	10	9	176	4,044	6,512	3,126	1,197	789	168	37	3	1	12
1950	14,930	9	3	5	9	10	7	7	12	6	11	205	3,917	5,913	2,883	1,037	698	142	33	9	------	14
1951	15,779	11	4	3	2	4	6	5	5	11	6	215	4,417	6,190	2,859	1,154	694	139	33	5	------	16
1952	15,895	11	1	6	12	11	5	4	3	4	6	218	4,569	6,250	2,850	1,061	736	98	26	3	2	19

TABLE 8C.—REPORTED UNTREATED CASES OF GONORRHEA BY YEAR OF REPORT, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Report)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	2,450	77.3	2,104	133.5	342	21.4	1,366	61.1	1,139	101.6	227	20.4	1,079	115.4	965	212.1	114	23.7	5
1931	2,298	71.7	1,980	124.2	318	19.7	1,269	56.0	1,052	92.6	217	19.2	1,029	109.5	928	202.9	101	20.9	----------
1932	2,130	65.2	1,858	114.5	272	16.6	1,164	50.3	988	85.2	176	15.2	963	101.3	868	187.4	95	19.5	3
1933	2,176	65.6	1,884	114.2	292	17.5	981	41.6	811	68.6	170	14.5	1,167	121.4	1,053	224.9	114	23.1	28
1934	2,164	64.4	1,779	106.5	380	22.5	1,113	46.5	867	72.3	246	20.6	1,041	107.6	908	192.4	133	26.8	10
1935	1,826	54.0	1,522	90.5	304	17.9	1,000	41.5	798	66.1	202	16.8	823	84.8	721	152.2	102	20.5	3
1936	2,159	63.5	1,531	90.6	628	36.7	1,159	47.7	749	61.6	410	33.8	996	102.3	779	163.9	217	43.6	4
1937	2,894	84.3	2,091	122.6	803	46.5	1,406	57.3	973	79.3	433	35.3	1,454	148.6	1,086	227.1	368	73.6	34
1938	3,045	87.6	2,401	139.0	644	36.8	1,252	50.3	939	75.4	313	25.1	1,748	177.5	1,432	297.2	316	62.8	45
1939	3,815	108.0	2,999	170.8	816	45.9	1,311	51.7	967	76.3	344	27.1	2,504	251.6	2,032	417.0	472	93.0	----------
1940	4,616	129.2	3,557	200.6	1,059	58.9	1,546	60.2	1,150	89.7	396	30.8	3,070	305.8	2,407	490.0	663	129.3	----------
1941	5,834	159.0	3,966	217.9	1,868	101.0	1,491	56.6	949	72.2	542	41.0	4,343	419.6	3,017	595.7	1,326	250.9	----------
1942	9,637	262.9	7,266	402.9	2,371	127.3	2,666	101.3	1,937	148.9	729	54.8	6,971	674.4	5,329	1,061.2	1,642	308.9	----------
1943	10,870	295.1	7,247	404.9	3,623	191.4	3,101	117.6	1,916	149.5	1,185	87.4	7,769	742.7	5,331	1,048.9	2,438	453.4	----------
1944	10,892	303.0	7,389	433.6	3,503	185.3	4,681	181.9	3,106	256.4	1,575	115.6	6,211	608.4	4,283	869.6	1,928	364.8	----------
1945	13,162	374.5	8,175	499.2	4,987	265.7	5,828	231.9	3,692	319.3	2,136	157.5	7,334	732.2	4,483	931.1	2,851	548.1	----------
1946	16,082	441.7	10,636	597.4	5,446	292.7	5,087	195.2	3,349	262.1	1,738	130.8	10,995	1,063.4	7,287	1,450.4	3,708	697.6	----------
1947	14,169	381.1	9,582	524.9	4,587	242.4	3,232	121.4	2,153	164.0	1,079	79.9	10,937	1,035.8	7,429	1,447.9	3,508	646.2	----------
1948	14,962	393.9	10,260	550.2	4,702	243.2	2,982	109.7	1,932	144.1	1,050	76.2	11,980	1,110.7	8,328	1,588.9	3,652	658.6	----------
1949	16,173	418.6	10,846	571.7	5,327	270.8	2,949	106.6	1,947	142.7	1,002	71.4	13,224	1,205.1	8,899	1,668.7	4,325	766.7	----------
1950	14,930	367.6	9,845	488.1	5,085	248.7	2,042	68.5	1,389	93.2	653	43.7	12,888	1,194.6	8,456	1,603.7	4,432	803.6	----------
1951	15,779	382.7	10,287	502.4	5,492	264.6	1,977	65.1	1,335	88.1	642	42.3	13,802	1,268.4	8,952	1,683.4	4,850	871.7	----------
1952	15,895	381.0	9,962	480.8	5,933	282.5	2,037	66.2	1,395	90.8	642	41.7	13,858	1,264.8	8,567	1,600.2	5,291	944.4	----------

TABLE 9A.—REPORTED UNTREATED CASES OF GRANULOMA INGUINALE BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF REPORT, 1952

(Place of Report)

Area	1951 Reports				1952 Reports																
	Color				Color				Month of report												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	58	1	57		38	1	37		5	5	2	5	1	4	4	1	1	2	6	2	
Alamance	1		1																		
Beaufort	2		2		1		1									1					
Bladen	1		1		1		1												1		
Cleveland	1		1																		
Columbus	4		4		2		2								1				1		
Craven	1		1																		
Cumberland	8	1	7																		
Duplin	3		3																		
Durham	3		3		3		3						1	1						1	
Forsyth					1		1		1												
Gates					2		2			1	1										
Guilford	2		2		1		1			1											
Johnston					1		1		1												
Lenoir					3		3			1								1	1		
Mecklenburg	16		16		10		10		3	1		3						1	1	1	

Nash					1		1										1				
New Hanover	6		6																		
Northampton	1		1																		
Perquimans	1		1																		
Pitt					1		1							1							
Richmond	2		2		1		1			1											
Robeson					7	1	6				1	2		1	1				2		
Rockingham	1		1																		
Vance	1		1																		
Wake	1		1		1		1								1						
Wilson	3		3		2		2							1	1						

TABLE 9B.—REPORTED UNTREATED CASES OF GRANULOMA INGUINALE BY YEAR OF REPORT AND AGE, 1945-1952
(Place of Report)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1945	75	------	------	------	------	------	------	------	------	------	------	------	25	15	15	8	5	5	2	------	------	------
1946	101	------	------	------	------	------	------	------	------	------	------	1	25	29	19	9	11	3	2	------	2	------
1947	99	------	------	------	------	------	------	------	------	------	------	1	26	29	21	6	11	2	1	2	------	------
1948	98	------	------	1	------	------	------	------	------	------	------	1	20	20	20	7	19	6	1	3	------	------
1949	66	------	------	------	------	------	------	------	------	------	------	------	8	19	10	8	14	4	------	2	------	1
1950	82	------	------	------	------	------	------	------	------	------	------	------	6	26	16	8	17	7	2	------	------	------
1951	58	------	------	------	------	------	------	------	------	------	------	1	5	21	9	8	9	2	1	1	------	1
1952	38	------	------	------	------	------	------	------	------	------	------	1	4	10	7	5	8	2	1	------	------	------

Table 9C.—Reported Untreated Cases of Granuloma Inguinale by Year of Report, Color, and Sex With Rates Per 100,000 Population, 1945-1952 (Place of Report)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1945	75	2.1	42	2.6	33	1.8	2	0.1	------	------	2	0.1	73	7.3	42	8.7	31	6.0	------
1946	101	2.8	51	2.9	50	2.7	5	0.2	4	0.3	1	0.1	96	9.3	47	9.4	49	9.2	------
1947	99	2.7	42	2.3	57	3.0	1	0.0	------	------	1	0.1	98	9.3	42	8.2	56	10.3	------
1948	98	2.6	52	2.8	46	2.4	3	0.1	3	0.2	------	------	95	8.8	49	9.3	46	8.3	------
1949	66	1.7	38	2.0	28	1.4	------	------	------	------	------	------	66	6.0	38	7.1	28	5.0	------
1950	82	2.0	50	2.5	32	1.6	------	------	------	------	------	------	82	7.6	50	9.5	32	5.8	------
1951	58	1.4	34	1.7	24	1.2	1	0.0	1	0.1	------	------	57	5.2	33	6.2	24	4.3	------
1952	38	0.9	15	0.7	23	1.1	1	0.0	------	------	1	0.1	37	3.4	15	2.8	22	3.9	------

TABLE 10A.—REPORTED CASES OF INFECTIOUS JAUNDICE BY COUNTY, COLOR, AND TYPE,* 1952 ONSETS
(Place of Residence)

Area	1952 Onsets						
	Total	White	Other	Infectious hepatitis		Serum hepatitis	
				White	Other	White	Other
North Carolina	675	625	50	622	50	3	
Alamance	4	4		4			
Alleghany	3	3		3			
Anson	1	1		1			
Ashe	42	42		42			
Avery	15	15		15			
Brunswick	1	1		1			
Buncombe	25	25		25			
Burke	8	8		8			
Cabarrus	2	2		2			
Caldwell	50	49	1	49	1		
Caswell	1	1		1			
Catawba	9	9		9			
Chatham	1	1		1			
Cherokee	3	3		3			
Clay	5	5		5			
Cleveland	22	21	1	21	1		
Craven	17	15	2	15	2		
Cumberland	23	19	4	19	4		
Dare	1	1		1			
Duplin	3	3		3			
Durham	31	28	3	28	3		
Edgecombe	5	3	2	3	2		
Franklin	1	1		1			
Gaston	22	22		22			
Graham	3	3		3			
Greene	3	3		3			
Guilford	6	6		5		1	
Halifax	20	15	5	15	5		
Haywood	48	47	1	47	1		
Jackson	22	18	4	18	4		
Jones	6	4	2	4	2		
Lenoir	10	9	1	9	1		
McDowell	1	1		1			
Macon	9	9		9			
Madison	1	1		1			

TABLE 10A.—JAUNDICE—*Continued*

Area	1952 Onsets						
	Total	White	Other	Infectious hepatitis		Serum hepatitis	
				White	Other	White	Other
Martin	5	5		5			
Mecklenburg	21	20	1	20	1		
Mitchell	21	21		21			
Montgomery	4	3	1	3	1		
Nash	12	11	1	11	1		
New Hanover	7	7		7			
Onslow	5	5		5			
Orange	13	13		13			
Pender	2	2		1		1	
Pitt	1	1		1			
Randolph	2	2		2			
Richmond	1	1		1			
Robeson	1	1		1			
Rowan	1	1		1			
Rutherford	2	2		2			
Sampson	24	23	1	23	1		
Stanly	3	3		2		1	
Stokes	8	5	3	5	3		
Surry	3	3		3			
Swain	14	5	9	5	9		
Vance	1		1		1		
Wake	2	2		2			
Watauga	3	3		3			
Wayne	6	6		6			
Wilson	16	9	7	9	7		
Yadkin	18	18		18			
Yancey	55	55		55			

* No cases of leptospirosis reported during year.

TABLE 10B.—REPORTED CASES OF INFECTIOUS JAUNDICE BY YEAR AND AGE, 1952
(Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1952	675	7	1	7	14	19	32	58	61	66	55	181	56	42	26	22	14	5	3	2	1	3

TABLE 10C.—REPORTED CASES OF INFECTIOUS JAUNDICE BY YEAR, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1952 (Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	Bo.	Rate	No.	Rate	
1952	675	16.2	366	17.7	309	14.7	625	20.3	334	21.7	291	18.9	50	4.6	32	6.0	18	3.2	----------

TABLE 11A.—REPORTED UNTREATED CASES OF LYMPHOGRANULOMA VENEREUM BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF REPORT, 1952
(Place of Report)

Area	1951 Reports				1952 Reports																
	Color				Color				Month of report												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	81	------	81	------	62	2	60	------	2	3	12	10	4	13	4	3	1	4	6	------	------
Alamance	2	------	2	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Anson	1	------	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Catawba	1	------	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Columbus	3	------	3	------	2	------	2	------	1	1	------	------	------	------	------	------	------	------	------	------	------
Craven	3	------	3	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Cumberland	10	------	10	------	20	1	19	------	------	------	6	------	1	8	2	------	------	1	2	------	------
Currituck	------	------	------	------	1	------	1	------	------	------	------	------	------	------	------	------	------	1	------	------	------
Duplin	2	------	2	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Durham	9	------	9	------	1	------	1	------	------	------	------	------	------	------	1	------	------	------	------	------	------
Edgecombe	2	------	2	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Forsyth	------	------	------	------	1	------	1	------	------	------	------	------	------	1	------	------	------	------	------	------	------
Franklin	------	------	------	------	1	------	1	------	------	------	------	------	------	1	------	------	------	------	------	------	------
Greene	2	------	2	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Guilford	2	------	2	------	5	------	5	------	------	------	1	2	------	2	------	------	------	------	------	------	------
Halifax	------	------	------	------	1	------	1	------	------	------	------	------	------	------	------	1	------	------	------	------	------
Harnett	------	------	------	------	1	------	1	------	------	------	------	1	------	------	------	------	------	------	------	------	------
Hoke	1	------	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Iredell	2	------	2	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Johnston	1	------	1	------	1	------	1	------	------	------	------	1	------	------	------	------	------	------	------	------	------
Lee	------	------	------	------	2	------	2	------	------	------	1	------	------	------	------	------	------	------	1	------	------

TABLE 11A.—LYMPHOGRANULOMA VENEREUM—*Continued*

Area	1951 Reports				1952 Reports																
	Color				Color				Month of report												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept	Oct.	Nov.	Dec.	Not stated
Lenoir	1		1																		
Martin					2		2		1	1											
Mecklenburg	4		4		1		1											1			
New Hanover	11		11		2		2				1		1								
Northampton	1		1																		
Onslow	3		3		1	1						1									
Orange					1		1												1		
Pasquotank					1		1				1										
Pitt	2		2		1		1						1								
Richmond	3		3																		
Robeson					7		7			1	1	3			1			1			
Rowan	1		1																		
Scotland					2		2						1				1				
Surry	1		1																		
Wake	1		1		1		1					1									
Washington	1		1																		
Wayne	2		2		3		3									2			1		
Wilson	9		9		4		4				1	1		1					1		

TABLE 11B.—REPORTED UNTREATED CASES OF LYMPHOGRANULOMA VENEREUM BY YEAR OF REPORT AND AGE, 1945-1952
(Place of Report)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1945	231	------	------	------	------	------	------	------	------	------	------	7	48	87	56	16	14	3	------	------	------	------
1946	153	------	------	------	------	------	------	------	------	------	------	3	34	66	37	6	5	2	------	------	------	------
1947	120	------	------	------	------	------	------	------	------	------	------	1	18	50	23	10	15	2	1	------	------	------
1948	176	------	------	------	------	------	------	------	------	------	------	1	17	59	51	19	24	5	------	------	------	------
1949	129	------	------	------	------	------	------	------	------	------	------	1	17	46	29	12	18	6	------	------	------	------
1950	91	------	------	------	------	------	------	------	------	------	------	1	9	33	21	10	13	2	2	------	------	------
1951	81	------	------	------	------	------	------	------	------	------	------	------	10	30	15	10	12	3	------	1	------	------
1952	62	------	------	------	------	------	------	------	------	------	------	1	9	18	9	8	10	6	------	------	------	1

TABLE 11C.—REPORTED UNTREATED CASES OF LYMPHOGRANULOMA VENEREUM BY YEAR OF REPORT, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1945-1952, (Place of Report)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1945	231	6.6	159	9.7	72	3.8	7	0.3	7	0.6	------	------	224	22.4	152	31.6	72	13.8	------
1946	153	4.2	114	6.4	39	2.1	5	0.2	5	0.4	------	------	148	14.3	109	21.7	39	7.3	------
1947	120	3.2	89	4.9	31	1.6	7	0.3	7	0.5	------	------	113	10.7	82	16.0	31	5.7	------
1948	176	4.6	132	7.1	44	2.3	4	0.1	3	0.2	1	0.1	172	15.9	129	24.6	43	7.8	------
1949	129	3.3	85	4.5	44	2.2	7	0.3	6	0.4	1	0.1	122	11.1	79	14.8	43	7.6	------
1950	91	2.2	70	3.5	21	1.0	3	0.1	3	0.2	------	------	88	8.2	67	12.7	21	3.8	------
1951	81	2.0	69	3.4	12	0.6	------	------	------	------	------	------	81	7.4	69	13.0	12	2.2	------
1952	62	1.5	46	2.2	16	0.8	2	0.1	2	0.1	------	------	60	5.5	44	8.2	16	2.9	------

Table 12A.—Reported Cases of Malaria by County and Color, 1951 and 1952, With Month of Onset, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																	
	Color				Color				Month of onset													
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	82	78	4	------	305	287	18	------	8	2	2	23	36	95	57	25	29	10	2	1	15
Brunswick	7	7	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Craven	1	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Edgecombe	2	1	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Mecklenburg	------	------	------	------	1	------	1	------	------	------	------	------	------	1	------	------	------	------	------	------	------
Nash	------	------	------	------	1	1	------	------	------	------	------	------	------	------	1	------	------	------	------	------	------
Onslow	2	2	------	------	1	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	1
Military sources	70	67	3	------	302	285	17	------	8	2	2	23	36	94	56	25	29	10	2	1	14

TABLE 12B.—REPORTED CASES OF MALARIA BY YEAR* AND AGE, 1937-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1937	876	16	8	21	20	14	17	12	13	18	19	57	77	90	85	59	118	94	49	21	13	55
1938	636	5	4	13	8	13	9	11	5	14	7	44	55	84	56	56	105	54	43	16	14	20
1939	629	5	13	12	7	19	17	11	10	4	9	49	34	55	82	61	90	49	49	30	8	15
1940	629	4	7	9	16	6	12	12	20	7	12	52	60	64	59	55	87	43	40	20	5	39
1941	237	------	2	7	------	5	4	4	5	4	1	10	24	31	24	17	40	22	16	14	------	7
1942	248	4	3	4	7	5	8	3	6	7	2	18	19	21	21	19	37	22	9	5	4	24
1943	183	2	2	6	------	3	1	1	1	2	2	5	15	26	12	14	24	13	11	9	2	32
1944	154	1	2	------	------	1	4	1	------	3	2	7	5	13	25	22	17	22	14	6	1	8
1945	554	------	2	12	2	2	1	8	4	4	5	16	22	72	122	68	74	68	47	19	3	3
1946	369	2	1	2	2	6	2	2	7	4	3	8	7	109	110	39	28	18	5	5	------	9
1947	137	1	------	9	3	4	3	1	3	2	2	5	------	23	24	15	18	11	5	3	------	5
1948	150	10	14	6	8	4	5	6	6	5	3	12	13	9	14	7	12	4	6	1	2	3
1949	53	------	1	------	1	3	3	2	4	1	1	3	8	8	5	2	5	5	------	------	------	1
1950	36	------	1	1	------	2	1	1	------	------	2	4	1	7	4	2	2	1	4	2	1	------
1951	82	------	------	1	2	------	1	------	------	------	------	1	8	39	11	11	4	1	------	3	------	------
1952	305	------	------	------	------	1	1	------	------	------	1	------	31	206	38	19	8	------	------	------	------	------

* 1937-1948 by year of report; 1949-1952 by year of onset.

Table 12C.—Reported Cases of Malaria by Year*, Color, and Sex With Rates per 100,000 Population, 1937-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1937	876	25.5	454	26.6	422	24.4	539	22.0	283	23.1	256	20.9	319	32.6	162	33.9	157	31.4	9
1938	636	18.3	293	17.0	343	19.6	425	17.1	199	16.0	226	18.2	200	20.3	90	18.7	110	21.9	11
1939	629	17.8	282	16.1	337	19.0	480	18.9	219	17.3	261	20.6	139	14.0	63	12.9	76	15.0	10
1940	629	17.6	286	16.1	343	19.1	442	17.2	204	15.9	238	18.5	187	18.6	82	16.7	105	20.5	----------
1941	237	6.5	108	5.9	129	7.0	167	6.3	81	6.2	86	6.5	70	6.8	27	5.3	43	8.1	----------
1942	248	6.8	106	5.9	141	7.6	200	7.6	86	6.6	114	8.6	47	4.5	20	4.0	27	5.1	1
1943	183	5.0	77	4.3	106	5.6	153	5.8	65	5.1	88	6.5	30	2.9	12	2.4	18	3.3	----------
1944	154	4.3	73	4.3	78	4.1	113	4.4	61	5.0	52	3.8	38	3.7	12	2.4	26	4.9	3
1945	554	15.8	312	19.1	240	12.8	405	16.1	269	23.3	136	10.0	147	14.7	43	8.9	104	20.0	2
1946	369	10.1	298	16.7	70	3.8	312	12.0	263	20.6	49	3.7	56	5.4	35	7.0	21	4.0	1
1947	137	3.7	83	4.5	54	2.9	101	3.8	64	4.9	37	2.7	36	3.4	19	3.7	17	3.1	----------
1948	150	3.9	78	4.2	72	3.7	118	4.3	63	4.7	55	4.0	32	3.0	15	2.9	17	3.1	----------
1949	53	1.4	34	1.8	19	1.0	46	1.7	29	2.1	17	1.2	7	0.6	5	0.9	2	0.4	----------
1950	36	0.9	21	1.0	15	0.7	30	1.0	17	1.1	13	0.9	6	0.6	4	0.8	2	0.4	----------
1951	82	2.0	74	3.6	8	0.4	78	2.6	70	4.6	8	0.5	4	0.4	4	0.8	------	------	----------
1952	305	7.3	305	14.7	------	------	287	9.3	287	18.7	------	------	18	1.6	18	3.4	------	------	----------

* 1937-1948 by year of report; 1949-1952 by year of onset.

TABLE 13A.—REPORTED CASES OF MEASLES BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	3,315	2,937	352	26	4,511	3,721	736	54	279	794	922	1,056	705	341	92	21	21	15	18	76	171
Alamance	19	19			38	37	1			6	13	10	7	2							
Alexander					14	14						7	4	3							
Alleghany					12	12					10	2									
Anson					2	2						1			1						
Ashe	8	8			566	557		9	2	307	256								1		
Avery					36	36				1	1	14	18		1						1
Beaufort	14	12	2		43	39	4		4	8	13	18									
Bertie	2	2			178	176	2				33	72	7								66
Bladen	2		2		2	2						1	1								
Brunswick	25	24	1		20	12	8				1	12	4	1	1					1	
Buncombe	112	110	1	1	224	182	41	1	16	46	66	71	21	4							
Burke	2	2			2	2									2						
Cabarrus	5	4	1		13	13					3	2	3	4							1
Caldwell					3	3					1	1			1						
Camden	1	1			1		1						1								

Carteret	143	139	2	2	13	13				3	1						8	1			
Caswell	65	47	7	11																	
Catawba	1	1			2		2					1						1			
Chatham	85	82	3		11	10	1			2	2		5	1					1		
Cherokee					3	3										3					
Chowan	4	2	2		3	3					3										
Clay					34	31	3					3	4	7	1				1	18	
Cleveland	1	1			28	28					4	9	13	2							
Columbus					1	1								1							
Craven	351	301	49	1	118	113	5			27	57	28	5			1					
Cumberland	24	23		1	20	17	3		2			5	2	1	1			1	1	7	
Currituck					8	4	4		1				1	2		1					3
Davidson	3	3			32	32					5	18	8	1							
Davie					3	3					3										
Duplin	1	1			16	15	1			4	1		1	6			2			2	
Durham	273	220	53		270	111	159		14	54	67	40	64	17	10		1	1	1		1
Edgecombe	59	49	10		132	123	8	1	59	46	19	3	4		1						
Forsyth	83	56	26	1	246	184	56	6	7	8	36	101	64	25	1		1		2	1	
Franklin	4	3	1		17	12	5		3	2	7	4	1								
Gaston	9	9			100	96	1	3	1	17	20	28	18	10	2	2			1		1
Gates	1	1			123		120	3		8	61	48				6					
Graham	1	1			21	21				2	6	12							1		
Granville	16	10	6		34	13	21		1		4	8	10	10	1						
Greene	1	1																			
Guilford	73	70	2	1	94	88		6	5		2	9	61	10	5					2	
Halifax	44	41	2	1	54	52	2		2	17	20	12	2		1						
Harnett	108	104	3	1	15	9	6			2	6	5			1				1		
Haywood	7	6	1		8	8				2	2	1	1	1		1					
Henderson	4	4			1		1						1								
Hertford	31	25	6		7	2	5		2		5										

TABLE 13A.—MEASLES—*Continued*

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Hoke	1	1			15		15			11	2		2								
Hyde	20	19	1																		
Iredell	5	5			22	21	1		1		2	1	7	10	1						
Jackson	3	1	1	1	105	104		1		1	6	48	29	5	2						14
Johnston	17	15	2		5	5			1		1		2								1
Jones	4	4			1		1									1					
Lee	41	33	8		11	11			5	2	3	1									
Lenoir	22	21	1		9	6	3				2	1	2	2			1	1			
Macon	1	1			306	305		1		3	9	201	27	8						1	57
Madison					19	19					4	14	1								
Martin	6	2	4																		
Mecklenburg	34	24	10		272	198	71	3	3	10	21	69	80	47	9		7	2		23	1
Mitchell	3	3			43	35		8		3	4	31	4			1					
Montgomery	8	8			6	5	1						6								
Moore	10	9	1		98	86	9	3	41	23	5	2	5	8	10	1	1				2
Nash	51	47	4		178	160	17	1	31	95	35	7	6	1							3
New Hanover	62	57	3	2	71	67	4		19	17	21	13	1								
Northampton	3		3		7	6	1			1		5	1								
Onslow	287	231	55	1	57	49	7	1	6	20	4							8	6	12	1
Orange	19	18	1		28	19	9		1	2	5	2	7	6	5						

Pamlico	25	17	8		1	1						1								
Pasquotank	12	11	1		23	4	19			2	1	3	7	8	2					
Pender					2	2					1		1							
Perquimans	12	12																		
Person	3	1	2		25	15	10				6	7	9	2						1
Pitt	181	142	38	1	19	15	4		4	6	4	5								
Polk	1		1		6	6					1	5								
Randolph	142	142			78	74	4				1	13	44	9	1			1	9	
Richmond	5	5			5	5					3		2							
Robeson	11	10	1		24	19	4	1		2	11	1	2	2	1			1		4
Rockingham	13	12	1		13	12		1				1	4	7						1
Rowan	5	5			1	1						1								
Rutherford					1	1					1									
Sampson	1	1																		
Stanly	5	3	2		49	46	3		30		1	8	7	3						
Stokes	1	1			22	22			1	2	13	3	3							
Surry	8	8			32	31	1					10	5	16	1					
Swain					18	18						1	13	4						
Transylvania	2	2																		
Tyrrell	4	4																		
Union					1	1						1								
Vance	3	3			10	5	5			5	3	2								
Wake	597	578	19		72	23	49		1		7	14	29	11	6	3				1
Warren	3	3			26	18	7	1	9	17										
Washington					7	7				3	3	1								
Watauga					175	171		4		1	1	3	59	77	22					12
Wayne	36	36			5	1	4					4	1							
Wilkes					17	17					1	13	3							
Wilson	61	55	5	1	47	20	27		5	4	8	17	4	7	1	1				
Yadkin	3	3			9	9			1	2	3	1	1		1					
Yancey	2	2			2	2			1		1									

TABLE 13B.—REPORTED CASES OF MEASLES BY YEAR* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	1,376	64	89	98	99	97	88	196	170	130	65	147	45	16	12	14	11	5	1	2	1	26
1931	15,499	280	582	758	922	1,026	974	2,008	1,945	1,566	1,133	2,530	829	317	174	99	95	47	14	5	5	190
1932	13,672	298	488	652	811	995	881	1,517	1,592	1,460	978	2,126	735	234	109	84	106	42	25	8	1	530
1933	16,197	304	492	734	808	1,004	1,103	1,946	1,894	1,984	1,452	2,965	565	262	98	96	73	41	17	9	3	347
1934	55,764	1,017	1,786	2,624	2,933	3,524	3,807	6,239	6,200	5,655	4,578	10,905	2,983	1,171	636	399	464	169	54	28	17	575
1935	10,683	260	417	529	586	688	642	1,094	1,047	952	742	2,259	654	237	105	59	65	26	7	5	1	308
1936	1,395	62	46	57	46	53	53	123	103	120	101	288	152	104	30	23	8	5	4	------	------	17
1937	7,937	188	274	332	407	439	437	757	854	933	712	1,516	539	206	77	49	68	24	5	3	------	117
1938	49,614	893	1,560	2,296	2,685	3,017	2,997	5,493	5,853	5,916	4,910	9,219	2,215	709	405	255	268	103	35	18	13	754
1939	20,031	381	661	821	1,082	1,242	1,232	2,223	2,544	2,143	1,763	4,088	801	260	160	119	121	60	80	8	2	240
1940	3,685	151	204	213	225	245	223	347	403	318	254	667	216	70	47	23	25	7	4	3	3	37
1941	27,764	601	963	1,479	1,874	1,895	1,979	3,354	3,449	2,919	1,891	4,459	1,186	464	286	196	160	60	27	7	2	513
1942	21,581	519	569	1,365	1,470	1,573	1,447	3,043	3,401	2,395	1,622	2,802	468	214	139	85	92	32	19	10	4	312
1943	5,217	207	152	373	387	420	386	593	638	434	235	524	278	176	99	61	49	14	6	1	1	183
1944	27,655	750	861	1,989	2,006	2,170	2,135	3,668	3,828	3,302	1,778	2,924	789	357	236	170	154	46	17	8	5	462
1945	1,194	63	50	126	104	102	104	181	134	93	45	101	46	14	13	6	2	------	------	------	------	10
1946	9,011	233	276	607	662	741	696	1,366	1,715	1,008	548	774	148	74	42	27	26	11	5	------	------	52
1947	4,978	156	159	317	367	422	429	932	821	580	271	325	61	28	30	19	12	4	2	1	------	42
1948	1,172	50	74	66	67	79	83	195	160	123	65	140	33	5	5	3	3	2	------	------	------	19
1949	17,330	363	727	909	934	1,036	1,117	2,922	3,045	2,095	1,112	2,194	274	80	45	47	59	11	5	2	4	349
1950	4,964	160	238	330	431	408	363	893	739	441	251	414	64	21	11	20	13	3	------	------	2	162
1951	3,315	77	107	155	211	239	194	544	560	368	177	301	122	193	25	7	12	2	2	1	------	18
1952	4,511	97	186	180	234	294	288	590	734	550	285	663	167	106	19	16	21	4	1	1	1	74

*1930–1948 by year of report; 1949–1952 by year of onset.

TABLE 13C.—REPORTED CASES OF MEASLES BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	1,376	43.4	695	44.1	670	42.0	1,240	55.5	634	56.6	601	53.9	125	13.4	54	11.9	68	14.2	11
1931	15,499	483.3	7,414	465.2	8,085	501.3	14,372	634.0	6,795	598.0	7,577	670.2	1,017	108.2	564	123.3	453	93.9	110
1932	13,672	418.8	6,462	398.2	6,781	413.0	11,097	479.6	5,478	472.3	5,619	486.9	2,051	215.7	955	206.2	1,096	224.7	524
1933	16,197	488.1	8,107	491.4	7,915	474.4	13,798	585.3	7,007	593.0	6,688	568.8	2,246	233.7	1,040	222.1	1,186	240.8	175
1934	55,764	1,660.0	27,775	1,662.6	27,836	1,648.4	48,372	2,022.3	24,215	2,020.0	24,033	2,014.3	6,923	715.6	3,306	700.7	3,588	724.0	469
1935	10,683	316.0	5,372	319.5	5,311	312.6	7,541	312.9	3,812	315.7	3,729	310.1	3,061	315.5	1,513	319.5	548	110.3	81
1936	1,395	41.0	713	42.2	682	39.9	1,229	50.6	640	52.7	589	48.6	154	15.8	66	13.9	88	17.7	12
1937	7,937	231.3	4,020	235.7	3,917	226.9	6,715	273.7	3,408	277.6	3,307	269.7	1,185	121.1	587	122.8	598	119.6	37
1938	49,614	1,427.5	24,668	1,428.0	24,946	1,427.1	42,299	1,698.2	21,284	1,708.7	21,015	1,687.8	7,007	711.5	3,204	665.0	3,803	756.1	308
1939	20,031	567.1	9,946	566.6	9,640	542.6	17,959	707.9	9,181	724.0	8,778	691.9	1,627	163.5	765	157.0	862	169.8	445
1940	3,685	103.2	1,873	105.6	1,810	100.6	2,604	101.4	1,337	104.3	1,267	98.5	1,078	107.4	535	108.9	543	105.9	3
1941	27,764	756.5	13,343	733.0	13,871	749.9	22,225	843.4	10,982	835.9	11,243	851.0	4,989	482.1	2,361	466.2	2,628	497.3	550
1942	21,581	588.8	10,610	588.3	10,441	560.7	18,599	706.7	9,367	719.9	9,232	693.9	2,452	237.2	1,243	247.5	1,209	227.5	530
1943	5,217	141.6	2,665	148.9	2,545	134.4	4,464	169.3	2,298	179.3	2,166	159.8	746	71.3	367	72.2	379	70.5	7
1944	27,655	769.3	13,642	800.6	13,479	712.8	24,797	963.4	12,437	1,026.6	12,360	907.2	2,324	227.6	1,205	244.7	1,119	211.8	534
1945	1,194	34.0	614	37.5	570	30.4	846	33.7	450	38.9	396	29.2	338	33.7	164	34.1	174	33.5	10
1946	9,011	247.5	4,527	254.3	4,443	238.8	7,263	278.6	3,685	288.4	3,578	269.3	1,707	165.1	842	167.6	865	162.7	41
1947	4,978	133.9	2,543	139.3	2,435	128.7	4,452	167.2	2,282	173.9	2,170	160.8	526	49.8	261	50.9	265	48.8	----------
1948	1,172	30.9	615	33.0	554	28.7	888	32.7	470	35.1	415	30.1	269	24.9	139	26.5	130	23.4	15
1949	17,330	448.5	8,756	461.5	8,556	435.0	14,106	509.9	7,180	526.4	6,910	492.6	3,125	284.8	1,534	287.7	1,591	282.0	99
1950	4,964	122.2	2,469	122.4	2,490	121.8	4,265	143.0	2,126	142.7	2,137	143.1	631	58.5	309	58.6	322	58.4	68
1951	3,315	80.4	1,799	87.9	1,514	72.9	2,937	96.8	1,614	106.5	1,323	87.1	352	32.3	175	32.9	177	31.8	26
1952	4,511	108.1	2,255	108.8	2,252	107.2	3,721	120.9	1,876	122.1	1,844	119.7	736	67.2	347	64.8	389	69.4	54

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 14A.—REPORTED CASES OF MENINGITIS BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	151	124	27		197	148	49		25	19	41	19	11	9	10	7	7	10	17	17	5
Alamance	1		1		3	2	1		1	1						1					
Alexander					1	1				1											
Ashe	1	1			2	2					1	1									
Avery					1	1				1											
Bertie					1		1				1										
Brunswick	3	2	1		1		1								1						
Buncombe	9	9			4	3	1		1			1				1				1	
Burke	6	5	1		5	5				1	1			2						1	
Cabarrus	2	1	1																		
Caldwell	4	4			3	3					2			1							
Carteret	1	1			2	2			1			1									
Caswell	1	1			2		2		1										1		
Catawba	2	1	1		1	1													1		
Chatham	1		1		2		2				1		1								
Cleveland	2	1	1		4	4			1	1	1						1				

Columbus	1	1			1		1			1											
Craven					1	1					1										
Cumberland	12	11	1		16	14	2		1	1	4	2	1	2	2	1		1			1
Currituck	1	1																			
Dare	1	1																			
Davidson	3	3			2	1	1				1	1									
Duplin					1		1				1										
Durham	1	1			3	3					1					1					1
Edgecombe	1	1			1		1												1		
Forsyth	5	5			6	6			2		1					1	1	1			
Franklin	4	1	3		1		1						1								
Gaston	2	1	1		2	2					1		1								
Greene	1	1			2	1	1					1							1		
Guilford	7	6	1		4	4					1	1	1							1	
Halifax	1	1			1		1				1										
Harnett	3	3			1	1							1								
Haywood					4	4				1	1						1		1		
Henderson	2	2			1	1											1				
Hertford	1		1		1		1												1		
Iredell	3	2	1		5	5					2	1						1		1	
Jackson					1	1									1						
Johnston	5	4	1		4	4				1	2	1									
Lee	2	2																			
Lenoir					3	3					1				1				1		
Lincoln					1	1													1		
McDowell					2	2					1								1		
Madison	1	1																			
Martin					3	2	1		1		1							1			
Mecklenburg	8	6	2		9	8	1		2	1	2					1		3			
Mitchell	4	4			5	5			1		3									1	

TABLE 14A.—MENINGITIS—*Continued*

Area	1951 Onsets				1952 Onsets															
	Color				Color				Month of onset											
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Montgomery	2	2			2	2													1	1	
Moore					1		1					1									
Nash	2	1	1		1	1												1			
New Hanover					1	1															1
Northampton	1		1		7	1	6		2	1	1					1			1	1	
Onslow	6	5	1		17	12	5		3	1	2	5			2		1	1	2		
Orange					2	1	1		1						1						
Pitt	1		1		5	2	3		1	2			1	1							
Polk					1	1									1						
Randolph	2	2																			
Robeson	3	1	2		5	3	2		1	1	1			1					1		
Rockingham					1	1											1				
Rowan	2	2			2	2			1										1		
Sampson	2	2			5	3	2				1		1							2	1
Scotland	1	1			3	2	1		1		1									1	
Stanly	1	1																			
Surry	4	4			2	2			1												1
Transylvania					1	1						1									
Union	1		1																		
Vance	2	1	1																		

Wake	9	8	1		10	7	3		1	1	1	1		1	1			1		3	
Warren	1	1																			
Washington					1	1							1								
Watauga					1	1														1	
Wayne	2	2			8	6	2			1	1	1					1		2	2	
Wilkes					1	1					1										
Wilson	3	3			4	1	3		1				2							1	
Yadkin	1	1			2	2				1				1							
Yancey	3	3			1	1				1											

TABLE 14B.—REPORTED CASES OF MENINGITIS BY YEAR* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	153	15	15	5	8	7	13	5	4	6	7	22	20	7	5	4	5	3	2	------	------	------
1931	123	10	8	6	6	7	5	4	7	2	3	21	22	7	2	6	5	------	------	------	------	2
1932	71	7	3	3	4	3	2	4	2	1	2	10	13	6	1	4	4	------	2	------	------	------
1933	66	14	4	5	2	1	4	4	3	------	------	9	10	3	1	1	1	4	------	------	------	------
1934	50	15	3	2	1	5	2	4	2	1	1	7	2	2	1	1	1	------	------	------	------	------
1935	109	14	6	3	1	4	2	3	1	2	3	17	16	13	10	4	2	7	1	------	------	------
1936	196	30	7	16	7	5	3	2	6	5	5	16	26	23	18	9	5	5	------	1	------	7
1937	169	19	5	9	6	7	7	4	7	2	6	19	27	17	8	5	10	7	2	------	------	2
1938	93	16	3	6	9	3	------	3	4	1	5	7	9	6	3	2	4	5	3	2	------	2
1939	59	10	4	6	2	3	2	2	2	1	------	4	7	6	2	3	2	------	1	------	------	2
1940	34	4	------	2	2	1	3	1	2	4	------	5	5	2	1	1	------	------	------	------	------	1
1941	46	7	2	1	1	------	2	3	------	------	2	9	7	5	2	------	1	2	------	------	------	2
1942	49	5	------	2	1	4	4	------	------	2	1	4	8	9	1	2	3	2	------	------	------	1
1943	478	19	2	14	4	7	5	8	7	4	1	26	122	114	63	30	28	6	2	1	------	15
1944	308	17	9	9	10	6	11	4	6	5	7	30	53	47	29	25	21	8	2	3	------	6
1945	179	11	8	8	4	8	15	4	2	5	1	14	38	16	10	8	15	8	1	------	------	3
1946	120	10	5	6	7	4	2	4	2	2	4	15	36	8	1	3	4	4	1	1	------	1
1947	95	17	2	10	6	1	5	2	5	6	3	10	6	5	1	2	4	5	2	------	1	2
1948	75	15	3	7	4	5	3	2	3	2	3	8	4	4	7	------	1	------	3	------	1	------
1949	87	9	7	7	7	4	4	4	2	1	3	12	9	6	1	2	------	5	2	1	1	------
1950	116	16	7	9	13	7	6	5	3	3	4	11	13	4	2	2	3	4	2	------	1	1
1951	151	12	17	13	13	11	7	8	4	5	2	14	15	5	2	4	5	10	1	3	------	------
1952	197	40	12	17	10	16	7	5	13	3	8	22	23	7	2	2	2	4	3	------	1	------

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 14C.—REPORTED CASES OF MENINGITIS BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	153	4.8	100	6.3	53	3.3	102	4.6	67	6.0	35	3.1	51	5.5	33	7.3	18	3.7	----------
1931	123	3.8	72	4.5	49	3.0	87	3.8	54	4.8	33	2.9	34	3.6	18	3.9	16	3.3	2
1932	71	2.2	43	2.6	28	1.7	45	1.9	23	2.0	22	1.9	26	2.7	20	4.3	6	1.2	----------
1933	66	2.0	38	2.3	28	1.7	44	1.9	25	2.1	19	1.6	22	2.3	13	2.8	9	1.8	----------
1934	50	1.5	35	2.1	15	0.9	36	1.5	26	2.2	10	0.8	14	1.4	9	1.9	5	1.0	----------
1935	109	3.2	77	4.6	32	1.9	78	3.2	51	4.2	27	2.2	31	3.2	26	5.5	5	1.0	----------
1936	196	5.8	137	8.1	59	3.5	124	5.1	86	7.1	38	3.1	72	7.4	51	10.7	21	4.2	----------
1937	169	4.9	120	7.0	49	2.8	83	3.4	60	4.9	23	1.9	85	8.7	59	12.3	26	5.2	1
1938	93	2.7	61	3.5	32	1.8	64	2.6	45	3.6	19	1.5	27	2.7	15	3.1	12	2.4	2
1939	59	1.7	45	2.6	14	0.8	45	1.8	34	2.7	11	0.9	14	1.4	11	2.3	3	0.6	----------
1940	34	1.0	25	1.4	9	0.5	24	0.9	20	1.6	4	0.3	10	1.0	5	1.0	5	1.0	----------
1941	46	1.3	31	1.7	15	0.8	35	1.3	24	1.8	11	0.8	11	1.1	7	1.4	4	0.8	----------
1942	49	1.3	37	2.1	10	0.5	33	1.3	27	2.1	6	0.5	14	1.4	10	2.0	4	0.8	2
1943	478	13.0	387	21.6	87	4.6	412	15.6	347	27.1	65	4.8	62	5.9	40	7.9	22	4.1	4
1944	308	8.6	229	13.4	74	3.9	258	10.0	194	16.0	64	4.7	45	4.4	35	7.1	10	1.9	5
1945	179	5.1	116	7.1	58	3.1	138	5.5	91	7.9	47	3.5	36	3.6	25	5.2	11	2.1	5
1946	120	3.3	86	4.8	33	1.8	95	3.6	71	5.6	24	1.8	24	2.3	15	3.0	9	1.7	1
1947	95	2.6	59	3.2	36	1.9	66	2.5	39	3.0	27	2.0	29	2.7	20	3.9	9	1.7	----------
1948	75	2.0	46	2.5	29	1.5	58	2.1	37	2.8	21	1.5	16	1.5	8	1.5	8	1.4	1
1949	87	2.3	40	2.1	47	2.4	69	2.5	31	2.3	38	2.7	18	1.6	9	1.7	9	1.6	----------
1950	116	2.9	57	2.8	59	2.9	84	2.8	41	2.8	43	2.9	32	3.0	16	3.0	16	2.9	----------
1951	151	3.7	78	3.8	73	3.5	124	4.1	61	4.0	63	4.1	27	2.5	17	3.2	10	1.8	----------
1952	197	4.7	110	5.3	87	4.1	148	4.8	86	5.6	62	4.0	49	4.5	24	4.5	25	4.5	----------

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 15A.—REPORTED CASES OF PARATYPHOID FEVER BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	5	4	1	------	4	4	------	------	------	------	1	------	------	------	2	------	1	------	------	------	------
Buncombe	1	------	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Caswell	------	------	------	------	2	2	------	------	------	------	------	------	------	------	2	------	------	------	------	------	------
Durham	1	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
Guilford	1	1	------	------	1	1	------	------	------	------	------	------	------	------	------	------	1	------	------	------	------
Robeson	------	------	------	------	1	1	------	------	------	------	1	------	------	------	------	------	------	------	------	------	------
Scotland	2	2	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

TABLE 15B.—REPORTED CASES OF PARATYPHOID FEVER BY YEAR* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	21									2	1	6	2	2	1		4	2	1			
1931	27								2	1	3	8	1	1	3	3	2	2	1			
1932	16				1	1			1	1	1	2	4	2		1		1	1			
1933	12						1			1	1	2	4	1	1		1					
1934	15	1								1		1	6	2			1	1		1		1
1935	26								1		1	2	1	3	6	4	5	3				
1936	20	1		1	1		1						6	1	1	4	1	1	1			1
1937	16			1					2				5	2	1	2		2			1	
1938	18	2			1				1	1		2	3	4	1		1	1	1			
1939	16	2			1	2				1		1	3	1			2	3				
1940	3						1			1		1										
1941	11	3				1							3		1	1		1	1			
1942	7		1	1				1		1		1					1		1			
1943	6						1			1		1	2						1			
1944	7	1										1	1		2			1	1			
1945	5									1				2	1					1		
1946	6	1		1									1				1	2				
1947	7			1								2	1	1			2					
1948	12		1	1			1	1		1		1		2		2	1			1		
1949	8										1	1	1		1		2		1		1	
1950	7											2	2	1			2					
1951	5								1					1	1			1		1		
1952	4		1								1	1		1								

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 15C.—REPORTED CASES OF PARATYPHOID FEVER BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	21	0.7	11	0.7	10	0.6	18	0.8	10	0.9	8	0.7	3	0.3	1	0.2	2	0.4	------
1931	27	0.8	17	1.1	10	0.6	21	0.9	16	1.4	5	0.4	6	0.6	1	0.2	5	1.0	------
1932	16	0.5	9	0.6	7	0.4	12	0.5	7	0.6	5	0.4	4	0.4	2	0.4	2	0.4	------
1933	12	0.4	10	0.6	2	0.1	9	0.4	8	0.7	1	0.1	3	0.3	2	0.4	1	0.2	------
1934	15	0.4	9	0.5	6	0.4	11	0.5	6	0.5	5	0.4	3	0.3	2	0.4	1	0.2	1
1935	26	0.8	17	1.0	9	0.5	16	0.7	9	0.7	7	0.6	10	1.0	8	1.7	2	0.4	------
1936	20	0.6	12	0.7	8	0.5	15	0.6	9	0.7	6	0.5	4	0.4	2	0.4	2	0.4	1
1937	16	0.5	7	0.4	9	0.5	9	0.4	6	0.5	3	0.2	7	0.7	1	0.2	6	1.2	------
1938	18	0.5	10	0.6	8	0.5	16	0.6	9	0.7	7	0.6	2	0.2	1	0.2	1	0.2	------
1939	16	0.5	8	0.5	7	0.4	14	0.6	8	0.6	6	0.5	1	0.1	------	------	1	0.2	1
1940	3	0.1	1	0.1	2	0.1	3	0.1	1	0.1	2	0.2	------	------	------	------	------	------	------
1941	11	0.3	4	0.2	7	0.4	7	0.3	2	0.2	5	0.4	4	0.4	2	0.4	2	0.4	------
1942	7	0.2	3	0.2	4	0.2	6	0.2	2	0.2	4	0.3	1	0.1	1	0.2	------	------	------
1943	6	0.2	3	0.2	3	0.2	4	0.2	1	0.1	3	0.2	2	0.2	2	0.4	------	------	------
1944	7	0.2	6	0.4	1	0.1	3	0.1	2	0.2	1	0.1	4	0.4	4	0.8	------	------	------
1945	5	0.1	1	0.1	4	0.2	4	0.2	1	0.1	3	0.2	1	0.1	------	------	1	0.2	------
1946	6	0.2	2	0.1	4	0.2	6	0.2	2	0.2	4	0.3	------	------	------	------	------	------	------
1947	7	0.2	4	0.2	3	0.2	5	0.2	3	0.2	2	0.1	2	0.2	1	0.2	1	0.2	------
1948	12	0.3	8	0.4	4	0.2	12	0.4	8	0.6	4	0.3	------	------	------	------	------	------	------
1949	8	0.2	5	0.3	3	0.2	8	0.3	5	0.4	3	0.2	------	------	------	------	------	------	------
1950	7	0.2	5	0.2	2	0.1	6	0.2	4	0.3	2	0.1	1	0.1	1	0.2	------	------	------
1951	5	0.1	3	0.1	2	0.1	4	0.1	2	0.1	2	0.1	1	0.1	1	0.2	------	------	------
1952	4	0.1	1	0.0	3	0.1	4	0.1	1	0.1	3	0.2	------	------	------	------	------	------	------

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 16A.—REPORTED CASES OF POLIOMYELITIS BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	314	287	27		538	498	40		12	5	2	2	8	37	91	150	112	83	21	15	
Alamance	3	3			9	5	4		1	1			1			2	2	1		1	
Alexander	5	5																			
Alleghany	3	3			2	2									1			1			
Ashe	4	4			8	8			3	2								1	2		
Avery	1	1																			
Beaufort					1	1									1						
Bertie	1		1		4	3	1							1		1		1	1		
Bladen	1		1		3	3										2	1				
Brunswick	2	1	1		3	3										1	1	1			
Buncombe	13	13			13	13								2	3	4	3	1			
Burke	13	10	3		23	22	1		1				1	1		17	1	2			
Cabarrus	2	2			4	4										2	1	1			
Caldwell	17	15	2		14	14			2						1	1		3	1	6	
Camden					1		1							1							
Carteret	2	2			4	4								1	1	2					

TABLE 16A.—POLIOMYELITIS—*Continued*

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Caswell	3	3																			
Catawba	18	18			8	8			1							3	1	2	1		
Chatham	1	1			2	1	1								1			1			
Cleveland	10	10			20	18	2			1	1		1	9	4	1	3				
Columbus	3	3			3	3									2	1					
Craven	1	1			6	6					1			1	1	1	1	1			
Cumberland	9	8	1		8	8						1				4		3			
Davidson	1	1			9	9			1						1	2	1	4			
Davie	1		1		4	4										3		1			
Duplin					1	1									1						
Durham	8	8			19	17	2								7	5	4	3			
Edgecombe	1		1		5	4	1									1	3	1			
Forsyth	2	2			41	41								1	1	15	19	5			
Franklin	2	1	1		2	1	1									1	1				
Gaston	10	10			19	19								2	8	5		2	1	1	
Granville	3	3																			
Greene					7	4	3						3	2	1	1					
Guilford	19	17	2		63	62	1			1				1	15	15	18	10	2	1	
Halifax					6	5	1										3	2		1	
Harnett	1	1			1	1										1					

Haywood	1	1			5	5								1	2			2			
Henderson	14	14			5	4	1								3	1	1				
Hoke	1	1			1	1										1					
Iredell	1	1			6	6											2	2	1	1	
Jackson	1	1			3	3											1	1	1		
Johnston	5	5			5	5						1		2		2					
Lee	5	5			4	4								1	1	1		1			
Lenoir					3	2	1								1	1	1				
McDowell	5	5			2	2										1	1				
Macon	3	3			3	3			1						1			1			
Madison					1	1										1					
Martin	1	1																			
Mecklenburg	15	15			9	7	2									2	4	2		1	
Mitchell	1	1																			
Montgomery					5	5			1								2	2			
Moore	2	2			4	4								1	2			1			
Nash	5	5			2	2											1			1	
New Hanover	6	5	1		10	8	2							1	1	2	4	2			
Northampton	1		1																		
Onslow	5	5			2	2									1	1					
Orange	12	12			9	9										3	2	2	2		
Pamlico					1	1												1			
Pasquotank	2	1	1																		
Person	1	1			1	1										1					
Pitt					7	5	2						1	1	4		1				
Polk	1	1			1	1										1					
Randolph	5	4	1		20	19	1								4	7	3	2	4		
Richmond					5	5							1		1		1	2			
Robeson	1	1			6	4	2								1		3		2		
Rockingham	7	7			29	27	2		1					2	4	9	9	4			

TABLE 16A.—POLIOMYELITIS—*Continued*

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Rowan	8	7	1		6	6											1	3	1	1	
Rutherford	4	4			4	3	1										1	3			
Sampson	1		1		4	4									2	2					
Scotland	1	1			2	1	1											1	1		
Stanly					4	4										3		1			
Stokes					2	2									1	1					
Surry	3	3			7	7								1	1	3	2				
Swain	1	1																			
Transylvania	2	1	1		10	8	2							2	2	4	1	1			
Union	4	4			3	3											2	1			
Vance	2	2			1	1									1						
Wake	12	8	4		13	13								1	4	6	2				
Warren					1	1									1						
Washington	2	2																			
Watauga	3	3			1	1										1					
Wayne	2	1	1		5	4	1							1	3	1					
Wilkes	4	3	1		5	3	2								1		2	1	1		
Wilson	4	4			3	2	1							1			1			1	
Yadkin					5	5										4	1				

TABLE 16B.—REPORTED CASES OF POLIOMYELITIS BY YEAR* AND AGE, 1931-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1931	104	11	29	17	13	4	2	4	3	3	5	5	3	1	1	1	1	------	------	------	------	1
1932	54	5	8	7	6	6	2	5	2	2	1	2	2	1	------	2	1	1	------	------	------	1
1933	30	2	6	2	2	3	3	2	1	1	------	5	------	------	1	------	------	------	------	------	------	2
1934	45	2	5	5	3	4	6	1	3	3	------	6	6	------	1	------	------	------	------	------	------	------
1935	675	52	68	78	85	53	52	38	33	30	35	62	36	15	8	7	10	5	3	1	------	4
1936	52	4	6	6	6	7	3	3	1	3	3	5	4	------	1	------	------	------	------	------	------	------
1937	106	10	11	21	10	8	9	5	5	4	2	14	2	2	2	------	------	------	------	------	------	1
1938	50	4	9	5	7	3	3	4	2	2	1	8	1	------	1	------	------	------	------	------	------	------
1939	116	5	15	10	9	13	9	10	8	2	7	14	6	3	1	2	1	------	------	------	------	1
1940	74	12	8	13	5	5	3	3	------	5	5	12	3	------	------	------	------	------	------	------	------	------
1941	168	6	10	31	20	16	9	13	12	14	6	22	5	1	1	------	------	------	------	------	------	2
1942	82	3	2	17	9	11	6	3	4	3	6	10	5	1	1	1	------	------	------	------	------	------
1943	36	5	2	4	6	7	2	------	------	2	------	4	3	------	1	------	------	------	------	------	------	------
1944	861	30	30	75	78	77	49	71	59	56	67	143	77	16	10	12	7	3	1	------	------	------
1945	159	7	9	21	12	9	15	12	10	7	5	29	14	4	4	1	------	------	------	------	------	------
1946	157	7	10	18	17	18	16	15	13	9	6	20	5	1	2	------	------	------	------	------	------	------
1947	300	12	20	40	41	25	27	16	21	12	14	47	13	7	3	------	1	------	1	------	------	------
1948	2,498	143	263	260	245	229	206	170	135	103	74	336	161	83	50	19	16	5	------	------	------	------
1949	229	9	17	23	22	17	24	20	13	5	8	30	15	10	8	4	3	1	------	------	------	------
1950	756	27	44	48	62	56	52	55	49	38	27	156	65	35	19	9	9	2	1	------	------	2
1951	314	16	18	20	22	24	21	19	17	19	24	54	23	22	8	3	4	------	------	------	------	------
1952	538	20	37	36	40	38	30	40	28	31	30	100	42	22	27	9	6	1	------	------	------	1

* 1931-1948 by year of report; 1949-1952 by year of onset.

TABLE 16C.—REPORTED CASES OF POLIOMYELITIS BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1931-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1931	104	3.2	64	4.0	40	2.5	88	3.9	54	4.8	34	3.0	14	1.5	8	1.7	6	1.2	2
1932	54	1.7	28	1.7	26	1.6	43	1.9	23	2.0	20	1.7	9	0.9	3	0.6	6	1.2	2
1933	30	0.9	15	0.9	15	0.9	19	0.8	7	0.6	12	1.0	10	1.0	7	1.5	3	0.6	1
1934	45	1.3	20	1.2	25	1.5	32	1.3	18	1.5	14	1.2	13	1.3	2	0.4	11	2.2	----------
1935	675	20.0	364	21.7	311	18.3	519	21.5	289	23.9	230	19.1	152	15.7	73	15.4	79	15.9	4
1936	52	1.5	28	1.7	24	1.4	39	1.6	23	1.9	16	1.3	13	1.3	5	1.1	8	1.6	----------
1937	106	3.1	68	4.0	38	2.2	91	3.7	58	4.7	33	2.7	15	1.5	10	2.1	5	1.0	----------
1938	50	1.4	22	1.3	28	1.6	40	1.6	17	1.4	23	1.8	10	1.0	5	1.0	5	1.0	----------
1939	116	3.3	48	2.7	68	3.8	95	3.7	40	3.2	55	4.3	21	2.1	8	1.6	13	2.6	----------
1940	74	2.1	49	2.8	25	1.4	61	2.4	40	3.1	21	1.6	13	1.3	9	1.8	4	0.8	----------
1941	168	4.6	81	4.4	79	4.3	146	5.5	74	5.6	72	5.4	14	1.4	7	1.4	7	1.3	8
1942	82	2.2	45	2.5	35	1.9	72	2.7	41	3.2	31	2.3	8	0.8	4	0.8	4	0.8	2
1943	36	1.0	24	1.3	12	0.6	30	1.1	20	1.6	10	0.7	6	0.6	4	0.8	2	0.4	----------
1944	861	24.0	466	27.3	395	20.9	786	30.5	424	35.0	362	26.6	75	7.3	42	8.5	33	6.2	----------
1945	159	4.5	86	5.3	73	3.9	135	5.4	76	6.6	59	4.3	24	2.4	10	2.1	14	2.7	----------
1946	157	4.3	87	4.9	69	3.7	131	5.0	69	5.4	62	4.7	25	2.4	18	3.6	7	1.3	1
1947	300	8.1	165	9.0	135	7.1	268	10.1	154	11.7	114	8.4	32	3.0	11	2.1	21	3.9	----------
1948	2,498	65.8	1,424	76.4	1,074	55.6	2,214	81.4	1,247	93.0	967	70.1	284	26.3	177	33.8	107	19.3	----------
1949	229	5.9	140	7.4	89	4.5	204	7.4	128	9.4	76	5.4	25	2.3	12	2.3	13	2.3	----------
1950	756	18.6	440	21.8	316	15.5	666	22.3	379	25.4	287	19.2	90	8.3	61	11.6	29	5.3	----------
1951	314	7.6	182	8.9	132	6.4	287	9.5	164	10.8	123	8.1	27	2.5	18	3.4	9	1.6	----------
1952	538	12.9	293	14.1	245	11.7	498	16.2	269	17.5	229	14.9	40	3.7	24	4.5	16	2.9	----------

* 1931-1948 by year of report; 1949-1952 by year of onset.

TABLE 17A.—REPORTED CASES OF ROCKY MOUNTAIN SPOTTED FEVER BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	61	57	4		38	29	8	1	1	1		2	5	11	9	3	5	1			
Alamance	1	1																			
Cabarrus					1	1								1							
Caswell	1	1																			
Chatham	6	5	1		1		1						1								
Cherokee	1	1																			
Cumberland					1	1											1				
Davidson	4	4			1	1										1					
Duplin					1	1							1								
Durham	2	2			2	2										1		1			
Edgecombe	1	1																			
Forsyth	1	1																			
Franklin					1	1									1						
Gaston	1	1																			
Granville					1		1							1							
Guilford	6	6			6	6								2	3	1					
Iredell	2	2																			
Lee	1	1																			
Martin	1	1																			

TABLE 17A.—ROCKY MOUNTAIN SPOTTED FEVER—*Continued*

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Mecklenburg	1	1			3	3							1	2							
Montgomery	2	2			1	1									1						
Moore	1	1																			
Nash	1	1			1		1								1						
Onslow	1	1			1	1			1												
Orange	4	2	2		2	1	1							1			1				
Person	2	1	1																		
Pitt	1	1																			
Randolph	5	5			1	1								1							
Richmond					1	1								1							
Robeson	1	1			7	2	4	1		1		1	2	2	1						
Rockingham	1	1																			
Rowan	3	3			2	2											2				
Rutherford	1	1			1	1											1				
Sampson	2	2			1	1									1						
Stanly	2	2			1	1									1						
Stokes	1	1																			
Surry	2	2																			
Warren	1	1																			
Wayne	1	1			1	1						1									

TABLE 17B.—REPORTED CASES OF ROCKY MOUNTAIN SPOTTED FEVER BY YEAR* AND AGE, 1932-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1932	1											1										
1933	27			1			1		2	3	1	4	4	3	1		2	2		2		1
1934	34		2		1		2	3		1	3	9	4	2	3		2		2			
1935	21				2		1		2	1		3	4	2		3	1	1	1			
1936	32			1	1		3	3	2	1	1	6	2	4	3	1		3	1			
1937	27			1	1	2	1	1	2	2		8	2	3	1		2		1			
1938	42		1	1	1	2	3	3	1	3		6	4	2	1	5	5	3	1			
1939	41			2	2	4	1	2	3	1	2	2	4	4	2	2		3	3	4		
1940	32				2	2	5	1				10	5	1			1	5				
1941	20			1		2	2	1	1	2		2	2			2	1	3	1			
1942	35			1	3	1	1	2	3	3	2	4		1	1	4	2	1	3	1		2
1943	36			2	1	1	1		5	4		10	2		2		2	2	2	2		
1944	59		3	2	6	1	3	4	5	1	4	11	10	1		3	5					
1945	57			5	4	2	3	5	6	3		6	1	5		4	4	3	4	2		
1946	66		3	4	3	3	7	4	4	6	3	7	5	4	1	2	5	1	2	1	1	
1947	88	1		4	5	4	2	4	9	7	5	15	6	2	1	3	7	8	1	1	3	
1948	74	2	2	2	8	6	3	5	2	8	3	12	1	1	1	4	7	2	3	1	1	
1949	79		3	1	4	5	5	6	5	5		13	2	3	8	4	3	4	3	3	1	1
1950	70	1	2	1	5	4	5	2	4	2	7	12	1	2	4	1	3	4	3	5	1	1
1951	61		1	4	5	8	3	1	4	2	5	7		2	1	2	3	7	4	2		
1952	38	1		2	1	4	3	3	1	2	2	7		4	3		1	2		1	1	

* 1932-1948 by year of report; 1949-1952 by year of onset.

TABLE 17C.—REPORTED CASES OF ROCKY MOUNTAIN SPOTTED FEVER BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1932-1952 (Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1932	1	0.03	1	0.06	------	------	1	0.04	1	0.08	------	------	------	------	------	------	------	------	------
1933	27	0.8	22	1.3	5	0.3	26	1.1	21	1.8	5	0.4	1	0.1	1	0.2	------	------	------
1934	34	1.0	18	1.1	16	0.9	31	1.3	16	1.3	15	1.3	1	0.1	1	0.2	------	------	2
1935	21	0.6	15	0.9	6	0.4	20	0.8	14	1.2	6	0.5	1	0.1	1	0.2	------	------	------
1936	32	0.9	19	1.1	13	0.8	31	1.3	18	1.5	13	1.1	1	0.1	1	0.2	------	------	------
1937	27	0.8	14	0.8	13	0.8	25	1.0	14	1.1	11	0.9	2	0.2	------	------	2	0.4	------
1938	42	1.2	25	1.4	17	1.0	37	1.5	22	1.8	15	1.2	4	0.4	2	0.4	2	0.4	1
1939	41	1.2	23	1.3	18	1.0	40	1.6	22	1.7	18	1.4	1	0.1	1	0.2	------	------	------
1940	32	0.9	15	0.8	17	0.9	29	1.1	13	1.0	16	1.2	3	0.3	2	0.4	1	0.2	------
1941	20	0.5	10	0.5	10	0.5	16	0.6	8	0.6	8	0.6	4	0.4	2	0.4	2	0.4	------
1942	35	1.0	23	1.3	11	0.6	29	1.1	19	1.5	10	0.8	5	0.5	4	0.8	1	0.2	1
1943	36	1.0	25	1.4	11	0.6	34	1.3	25	2.0	9	0.7	2	0.2	------	------	2	0.4	------
1944	59	1.6	34	2.0	24	1.3	57	2.2	33	2.7	24	1.8	1	0.1	1	0.2	------	------	1
1945	57	1.6	27	1.6	30	1.6	49	2.0	22	1.9	27	2.0	8	0.8	5	1.0	3	0.6	------
1946	66	1.8	28	1.6	38	2.0	56	2.1	22	1.7	34	2.6	10	1.0	6	1.2	4	0.8	------
1947	88	2.4	45	2.5	43	2.3	79	3.0	38	2.9	41	3.0	9	0.9	7	1.4	2	0.4	------
1948	74	1.9	39	2.1	35	1.8	66	2.4	35	2.6	31	2.2	7	0.6	3	0.6	4	0.7	1
1949	79	2.0	44	2.3	35	1.8	73	2.6	38	2.8	35	2.5	6	0.5	6	1.1	------	------	------
1950	70	1.7	44	2.2	26	1.3	59	2.0	37	2.5	22	1.5	11	1.0	7	1.3	4	0.7	------
1951	61	1.5	30	1.5	31	1.5	57	1.9	28	1.8	29	1.9	4	0.4	2	0.4	2	0.4	------
1952	38	0.9	16	0.8	22	1.0	29	0.9	12	0.8	17	1.1	8	0.7	4	0.7	4	0.7	1

* 1932-1948 by year of report; 1949-1952 by year of onset.

TABLE 18A.—REPORTED CASES OF SCARLET FEVER BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	2,236	2,132	94	10	2,222	2,115	101	6	313	375	303	184	75	46	32	39	116	264	247	220	8
Alamance	60	58	2		92	91	1		12	10	27	24	9	1				2		7	
Alexander	16	16			34	34			3	1	1	1					4	5	13	6	
Alleghany	3	3			2	2			2												
Anson	11	10	1		5	3	2						1				2	2			
Ashe	18	18			65	65			4	28	28	1		1			1		2		
Avery	2	2			8	8			3	2								1	2		
Beaufort	9	9			2	2			1									1			
Bertie	1	1			2	1	1											1	1		
Bladen					6	6				1							1	3		1	
Brunswick	2	2			3	3				2				1							
Buncombe	75	66	9		58	52	5	1	10	7	3	5	2	1	2	1	8	8	6	5	
Burke	78	75	3		50	49	1		13	15	1	1	1	3				10	6		
Cabarrus	45	43	1	1	67	66	1		25	8	12	3	2	1			3	5	5	3	
Caldwell	43	42	1		45	43	2		4	5	8	5	1			2	4	6	4	6	
Carteret					2	2			1	1											

TABLE 18A.—SCARLET FEVER—*Continued*

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Caswell	1		1		4	3	1			1								1	2		
Catawba	57	55	2		34	32	2		3	9	4	4	1			1	6	2	2	2	
Chatham	6	5	1		5	4	1								1		1	1	1		1
Cherokee	3	3																			
Chowan	6	6			1	1													1		
Clay					2	2			1											1	
Cleveland	45	45			48	40	8		6	6	3	7					5	5	2	14	
Columbus	8	8			3	1	2			1								1	1		
Craven	36	36			26	25	1		10	5	3			1				1	5	1	
Cumberland	14	14			67	65	2		4	2	5	1		1	2	3	7	25	5	11	1
Currituck					1	1			1												
Dare					1	1					1										
Davidson	50	50			75	74	1		16	13	3	10	5	1	1	2	4	3	10	7	
Davie	5	5			31	31				6	17	6	2								
Duplin					3	3													3		
Durham	81	76	5		143	136	7		14	17	12	16	6		1	2	11	21	28	15	
Edgecombe	25	22	3		16	11	5		2	2	1		3					2	4	2	
Forsyth	236	227	8	1	102	93	9		9	18	11	9	2	1		4	6	11	7	24	
Franklin	6	5	1		10	9	1		1	1									5	3	
Gaston	124	122	2		30	30			2	5	4	2			1	3	1	4	3	5	

Gates	2	2			1	1				1											
Graham	1	1																			
Granville	8	7	1		5	5							1					1		3	
Greene	2	2			5	5			3										2		
Guilford	184	172	9	3	132	126	5	1	16	22	34	14	13	2	3		3	9	10	6	
Halifax	11	11			6	4	2				1						1	2		2	
Harnett	15	15			15	14	1		1			3	6				1	3		1	
Haywood	30	30			21	19	1	1	7	1	5	1				1	1	1	1	3	
Henderson	19	18	1		14	14			3	8		1						2			
Hoke	4	1	3		9	9					2					3		1		3	
Hyde	1	1																			
Iredell	51	51			33	32	1		10	10	5		1	1	1			1	3	1	
Jackson	4	4			9	9			2	5	1							1			
Johnston	16	15	1		6	6			1			1						2		2	
Jones	3	3			2	2					1						1				
Lee	3	2	1		9	9				3						1		3	1	1	
Lenoir	9	9			12	10	1	1	1	2		1			2	1		2		3	
Lincoln	11	10	1		3	3			2							1					
Macon	1	1			36	36											8	8	11	7	2
Madison	4	4			4	4			1									2		1	
Martin	5	4		1	7	4	3		1				1		2	2	1				
Mecklenburg	196	195		1	129	128	1		16	16	13	19	4	10	2	1	6	12	21	9	
Mitchell	10	10			14	14				3		1	1	1					5		3
Montgomery	7	7			6	6						2		2	2						
Moore	5	5			2	1	1			1		1									
Nash	21	20	1		14	13	1		1	1				1		1	4	1	1	4	
New Hanover	32	31	1		64	62	1	1	6	8	19	7	1	1	1		5	7	5	4	
Northampton	3	1	1	1	5	3	2						1				1		2	1	
Onslow	10	6	4		14	14			1	2							1	2	6	2	
Orange	12	10	2		11	11				4	2		1					2		2	

TABLE 18A.—SCARLET FEVER—*Continued*

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Pamlico	2	2																			
Pender					1	1												1			
Perquimans					4	2	2		2	1									1		
Person	1	1			3	1	2				1									2	
Pitt	18	18			16	11	5		2		4	1		1		1	2	3	1	1	
Polk	2	2																			
Randolph	43	42		1	37	35	2		4	4	4	2	3	4			2	6	6	2	
Richmond	8	8			9	9				1					1	1	1	2	2	1	
Robeson	8	5	3		21	17	4		2	5	1	1	1	1			1	1	6	2	
Rockingham	7	6	1		10	9	1		1	2		3			1			1		2	
Rowan	42	41		1	58	56	2		12	8	11	4		1	1		4	11	3	3	
Rutherford	23	21	2		5	5				1	1					3					
Sampson	12	12			11	11					1				1		2	4	1	2	
Scotland	5	3	2		1	1									1						
Stanly	48	47	1		79	79			20	10	9	2		2			1	7	19	9	
Stokes	8	8			4	4			3	1											
Surry	50	50			83	82		1	13	39	7	3	1	1	1	2		11	4	1	
Swain	2	2																			
Transylvania	1	1			1	1												1			
Union	3	2	1																		

Vance	11	11			7	6	1		2		1						1	2	1		
Wake	67	55	12		129	123	6		16	16	24	17	4	3	5	2	4	12	11	15	
Warren	7	5	2		2	2													1	1	
Washington	1	1																			
Watauga	27	27			5	5			1	2			1	1							
Wayne	8	8			11	11								1				7		3	
Wilkes	20	20			16	16				1	1	3						2	2	7	
Wilson	14	12	2		16	13	3		3	2		2		1		1		5	1		1
Yadkin	38	36	2		53	53			13	29	11										
Yancey	14	14			9	9											1	5	2	1	

TABLE 18B.—REPORTED CASES OF SCARLET FEVER BY YEAR* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	3,096	63	104	202	275	311	321	374	297	277	189	354	88	45	32	15	9	2	1	1	------	136
1931	3,461	61	110	190	246	331	314	389	357	290	182	586	168	84	45	23	24	6	3	1	1	50
1932	2,868	57	86	167	197	278	227	324	277	277	193	492	131	65	35	21	21	7	2	------	------	11
1933	3,662	65	117	219	268	333	336	387	362	291	263	647	130	67	54	36	22	5	2	------	------	58
1934	2,852	41	106	172	235	275	265	314	253	261	153	485	118	67	33	25	11	7	2	2	------	27
1935	2,072	31	83	112	202	196	197	254	199	157	117	320	72	40	35	17	10	4	1	1	------	24
1936	1,921	36	62	107	155	196	166	196	193	169	117	335	77	43	28	16	8	2	------	1	------	14
1937	2,073	50	82	108	165	181	172	237	222	163	136	339	87	41	39	15	7	3	------	------	------	26
1938	2,228	41	82	135	165	204	179	223	210	195	166	369	126	40	32	22	11	2	1	1	------	24
1939	2,599	49	81	159	187	240	231	283	255	214	182	452	96	39	34	26	21	5	8	1	------	36
1940	2,442	42	74	152	196	240	235	336	235	217	143	364	81	50	17	16	12	2	1	------	------	29
1941	2,165	30	52	126	166	217	225	262	247	191	120	333	73	40	22	15	15	3	2	------	------	26
1942	2,570	49	40	145	221	266	262	326	251	218	173	411	79	35	20	31	18	3	2	------	------	20
1943	2,854	52	59	180	266	269	263	339	276	246	177	432	117	80	33	12	14	4	3	2	------	30
1944	2,239	35	55	147	209	214	203	257	224	175	141	324	85	61	38	24	13	2	2	1	------	29
1945	3,248	43	62	210	280	304	277	390	352	298	243	536	109	47	34	26	19	4	1	------	------	13
1946	1,530	32	24	85	134	147	141	181	138	124	91	223	117	48	21	9	9	------	3	1	------	2
1947	1,167	21	22	73	110	114	123	162	129	91	79	176	35	12	7	4	6	1	------	------	------	2
1948	1,099	14	46	79	106	126	123	169	129	94	50	132	12	7	3	3	3	------	------	------	------	3
1949	1,793	18	60	115	132	184	181	284	233	180	108	232	35	11	4	4	6	2	------	------	------	4
1950	1,885	21	55	119	200	192	182	263	216	198	143	226	34	9	6	9	8	1	------	------	------	3
1951	2,236	15	40	118	219	250	238	336	286	216	147	302	33	13	4	8	2	------	1	------	------	8
1952	2,222	18	52	107	181	229	248	356	265	208	169	314	40	6	9	7	4	3	1	------	------	5

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 18C.—REPORTED CASES OF SCARLET FEVER BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	3,096	97.6	1,295	82.2	1,674	104.9	2,882	128.9	1,249	111.4	1,603	143.7	79	8.4	31	6.8	48	10.0	137
1931	3,461	107.9	1,362	85.5	2,089	129.5	3,308	145.9	1,301	114.5	2,002	177.1	116	12.3	45	9.8	71	14.7	37
1932	2,868	87.8	1,238	76.3	1,616	98.4	2,774	119.9	1,204	103.8	1,570	136.0	69	7.3	29	6.3	40	8.2	25
1933	3,662	110.4	1,550	93.9	2,108	126.3	3,488	148.0	1,477	125.0	2,007	170.7	145	15.1	56	12.0	89	18.1	29
1934	2,852	84.9	1,174	70.3	1,673	99.1	2,737	114.4	1,127	94.0	1,605	134.5	100	10.3	40	8.5	60	12.1	15
1935	2,072	61.3	865	51.5	1,207	71.0	1,975	82.0	819	67.8	1,156	96.1	89	9.2	44	9.3	45	9.1	8
1936	1,921	56.5	804	47.6	1,117	65.3	1,865	76.8	786	64.7	1,079	89.0	52	5.3	17	3.6	35	7.0	4
1937	2,073	60.4	826	48.4	1,247	72.2	1,984	80.9	791	64.4	1,193	97.3	80	8.2	30	6.3	50	10.0	9
1938	2,228	64.1	960	55.6	1,268	72.5	2,129	85.5	916	73.5	1,213	97.4	91	9.2	40	8.3	51	10.1	8
1939	2,599	73.6	1,062	60.5	1,467	82.6	2,437	96.1	1,016	80.1	1,421	112.0	92	9.2	46	9.4	46	9.1	70
1940	2,442	68.4	1,100	62.0	1,342	74.6	2,240	87.2	1,011	78.9	1,229	95.6	202	20.1	89	18.1	113	22.0	----------
1941	2,165	59.0	913	50.2	1,204	65.1	1,962	74.5	839	63.9	1,123	85.0	155	15.0	74	14.6	81	15.3	48
1942	2,570	70.1	1,089	60.4	1,427	76.6	2,369	90.0	1,027	78.9	1,342	100.9	147	14.2	62	12.3	85	16.0	54
1943	2,854	77.5	1,277	71.3	1,569	82.9	2,643	100.2	1,187	92.6	1,456	107.4	236	22.6	90	17.7	113	21.0	8
1944	2,239	62.3	981	57.6	1,213	64.1	2,069	80.4	934	77.1	1,135	83.3	125	12.2	47	9.5	78	14.8	45
1945	3,248	92.4	1,390	84.9	1,817	96.8	3,070	122.2	1,329	114.9	1,741	128.3	137	13.7	61	12.7	76	14.6	41
1946	1,530	42.0	749	42.1	765	41.1	1,435	55.1	704	55.1	731	55.0	79	7.6	45	9.0	34	6.4	16
1947	1,167	31.4	560	30.7	607	32.1	1,108	41.6	532	40.5	576	42.7	59	5.6	28	5.5	31	5.7	----------
1948	1,099	28.9	494	26.5	605	31.3	1,041	38.3	473	35.3	568	41.2	48	4.5	18	3.4	30	5.4	10
1949	1,793	46.4	818	43.1	975	49.6	1,722	62.2	795	58.3	927	66.1	64	5.8	21	3.9	43	7.6	7
1950	1,885	46.4	870	43.1	1,015	49.6	1,787	59.9	832	55.8	955	64.0	93	8.6	38	7.2	55	10.0	5
1951	2,236	54.2	1,010	49.3	1,226	59.1	2,132	70.2	965	63.7	1,167	76.8	94	8.6	38	7.1	56	10.1	10
1952	2,222	53.3	1,015	49.0	1,207	57.5	2,115	68.7	966	62.9	1,149	74.6	101	9.2	45	8.4	56	10.0	6

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 19A.—REPORTED CASES OF SEPTIC SORE THROAT BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	29	28	1		33	30	3		6	5	3	1		2	3		1	9	2	1	
Ashe					3	3					3										
Bertie	2	2																			
Brunswick					1	1												1			
Buncombe	1	1																			
Cabarrus					1	1								1							
Catawba	1	1																			
Chatham					1		1					1									
Cumberland					1		1												1		
Dare					3	3											1	2			
Forsyth	1	1																			
Gaston	2	2																			
Halifax	7	7			5	5								1				4			
Jackson					5	5			2	3											
Johnston					1		1		1												
Mecklenburg	2	1	1		3	3			1						2						

Mitchell	1	1																			
Moore	1	1																			
Nash	1	1																			
Northampton					1	1									1						
Onslow	1	1																			
Stanly					3	3			2										1		
Swain					2	2				2											
Yadkin					1	1														1	
Yancey	9	9			2	2												2			

TABLE 19B.—REPORTED CASES OF SEPTIC SORE THROAT BY YEAR* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	114	2	1	4	8	6	2	8	6	5	2	16	9	10	9	10	5	4	2	1	1	3
1931	153	------	2	5	6	5	5	6	8	6	8	22	18	9	11	15	11	3	1	2	1	9
1932	142	------	1	4	5	2	7	3	3	5	1	18	23	24	10	12	11	6	4	------	------	3
1933	123	7	3	1	3	3	1	3	8	2	6	15	12	17	12	14	6	6	3	1	------	------
1934	104	------	1	2	7	7	7	3	4	4	1	15	6	17	6	5	7	4	1	1	------	6
1935	116	1	3	9	7	5	4	4	2	3	2	10	19	10	11	6	7	5	2	2	1	3
1936	115	3	5	2	5	6	2	11	10	1	4	12	8	6	14	8	7	4	1	2	1	3
1937	125	------	3	7	6	6	1	4	8	6	3	20	9	13	10	9	8	7	2	1	------	2
1938	148	2	4	5	8	6	6	4	8	3	3	25	19	18	7	7	11	5	1	1	------	5
1939	116	4	5	7	5	3	6	5	4	5	5	20	12	9	8	6	6	2	2	------	------	2
1940	88	------	4	2	4	1	2	3	3	1	4	9	12	13	5	4	7	7	------	5	------	2
1941	139	1	2	4	3	6	2	9	6	5	3	24	16	13	16	7	9	3	2	3	3	2
1942	109	5	2	5	5	2	3	4	5	5	1	13	17	10	6	5	12	5	2	------	------	2
1943	201	1	------	4	2	1	4	2	7	2	5	12	57	51	20	10	8	8	2	2	------	3
1944	47	------	------	1	1	2	2	1	------	2	4	9	12	3	4	------	2	2	1	1	------	------
1945	101	1	1	4	4	3	5	3	4	1	2	13	15	9	11	6	9	5	2	2	------	1
1946	51	1	------	4	5	4	1	2	3	------	3	1	8	4	5	1	4	2	2	------	------	1
1947	44	3	2	1	1	2	2	2	------	1	4	8	2	4	5	3	1	2	------	------	------	1
1948	30	1	3	1	1	2	1	1	1	------	------	6	2	2	3	2	1	1	1	------	1	------
1949	45	------	1	5	4	------	1	2	------	------	2	11	4	6	5	1	1	2	------	------	------	------
1950	41	1	1	------	2	1	------	1	1	2	------	7	4	4	6	5	4	2	------	------	------	------
1951	29	------	------	2	3	2	------	1	3	2	1	3	3	2	2	1	2	------	1	------	1	------
1952	33	2	1	2	2	1	1	2	3	2	1	2	3	1	3	2	3	------	2	------	------	------

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 19C.—REPORTED CASES OF SEPTIC SORE THROAT BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	114	3.6	49	3.1	65	4.1	92	4.1	40	3.6	52	4.7	17	1.8	8	1.8	9	1.9	5
1931	153	4.8	66	4.1	87	5.4	139	6.1	59	5.2	80	7.1	10	1.1	4	0.9	6	1.2	4
1932	142	4.3	60	3.7	82	5.0	127	5.5	53	4.6	74	6.4	14	1.5	6	1.3	8	1.6	1
1933	123	3.7	58	3.5	65	3.9	106	4.5	50	4.2	56	4.8	17	1.8	8	1.7	9	1.8	----------
1934	104	3.1	41	2.5	63	3.7	81	3.4	32	2.7	49	4.1	23	2.4	9	1.9	14	2.8	----------
1935	116	3.4	59	3.5	57	3.4	103	4.3	52	4.3	51	4.2	13	1.3	7	1.5	6	1.2	----------
1936	115	3.4	48	2.8	67	3.9	100	4.1	44	3.6	56	4.6	14	1.4	4	0.8	10	2.0	1
1937	125	3.6	59	3.5	66	3.8	94	3.8	48	3.9	46	3.8	29	3.0	11	2.3	18	3.6	2
1938	148	4.3	68	3.9	80	4.6	130	5.2	58	4.7	72	5.8	17	1.7	9	1.9	8	1.6	1
1939	116	3.3	53	3.0	61	3.4	106	4.2	48	3.8	58	4.6	8	0.8	5	1.0	3	0.6	2
1940	88	2.5	43	2.4	45	2.5	76	3.0	38	3.0	38	3.0	12	1.2	5	1.0	7	1.4	----------
1941	139	3.8	71	3.9	65	3.5	118	4.5	65	4.9	53	4.0	18	1.7	6	1.2	12	2.3	3
1942	109	3.0	51	2.8	54	2.9	93	3.5	43	3.3	50	3.8	12	1.2	8	1.6	4	0.8	4
1943	201	5.5	152	8.5	49	2.6	199	7.5	150	11.7	49	3.6	2	0.2	2	0.4	-------	-------	----------
1944	47	1.3	25	1.5	22	1.2	42	1.6	22	1.8	20	1.5	5	0.5	3	0.6	2	0.4	----------
1945	101	2.9	44	2.7	56	3.0	82	3.3	35	3.0	47	3.5	18	1.8	9	1.9	9	1.7	1
1946	51	1.4	22	1.2	29	1.6	41	1.6	18	1.4	23	1.7	10	1.0	4	0.8	6	1.1	----------
1947	44	1.2	19	1.0	25	1.3	39	1.5	16	1.2	23	1.7	5	0.5	3	0.6	2	0.4	----------
1948	30	0.8	13	0.7	17	0.9	26	1.0	12	0.9	14	1.0	4	0.4	1	0.2	3	0.5	----------
1949	45	1.2	22	1.2	23	1.2	43	1.6	21	1.5	22	1.6	1	0.1	1	0.2	-------	-------	1
1950	41	1.0	18	0.9	23	1.1	40	1.3	18	1.2	22	1.5	1	0.1	-------	-------	1	0.2	----------
1951	29	0.7	12	0.6	17	0.8	28	0.9	12	0.8	16	1.1	1	0.1	-------	-------	1	0.2	----------
1952	33	0.8	12	0.6	21	1.0	30	1.0	11	0.7	19	1.2	3	0.3	1	0.2	2	0.4	----------

* 1930-1948 by year of report; 1949-1951 by year of onset.

TABLE 20A.—REPORTED CASES OF SMALLPOX BY YEAR OF REPORT* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	556	6	6	4	6	8	6	10	5	6	6	25	68	89	52	45	74	68	35	16	3	18
1931	63	3	1	2	1	1	4	5	2	1	------	3	4	7	4	5	9	4	3	1	1	2
1932	72	1	------	1	3	2	3	2	3	1	1	5	9	7	9	5	11	4	3	------	------	2
1933	35	1	------	1	1	1	2	3	1	1	1	2	1	10	5	1	1	3	------	------	------	------
1934	18	1	------	------	------	------	------	------	------	------	------	3	2	5	2	1	3	1	------	------	------	------
1935	24	1	------	------	2	------	------	1	4	------	------	------	3	2	4	1	3	3	------	------	------	------
1936	19	------	1	------	------	------	------	------	------	------	------	2	------	5	2	3	4	------	1	------	------	1
1937	11	------	------	------	------	------	------	------	------	1	2	1	------	------	3	------	2	2	------	------	------	------
1938	35	------	------	1	------	------	------	------	------	1	1	1	------	3	5	5	12	2	2	1	1	------
1939	15	------	1	------	------	1	------	------	2	------	------	1	2	2	1	1	3	------	------	------	------	1
1940	6	------	------	------	1	1	------	------	------	------	------	------	------	1	------	------	1	2	------	------	------	------
1941	4	1	1	------	------	------	------	------	------	------	------	------	------	------	1	1	------	------	------	------	------	------
1942	7	------	------	------	------	------	------	------	------	------	------	2	------	2	------	------	3	------	------	------	------	------
1943	21	------	------	1	------	------	------	------	------	------	------	1	3	1	1	2	2	3	4	2	1	------
1944	4	------	------	1	2	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
1945	1	------	------	------	------	------	------	------	------	------	------	------	------	------	------	1	------	------	------	------	------	------
1946	2	------	------	------	------	------	------	------	------	------	------	1	------	------	------	------	------	1	------	------	------	------
1947	2	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	1	------	------	1	------
1948	3	------	------	------	------	------	------	------	1	------	------	------	------	1	------	------	------	------	1	------	------	------
1951	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
1952	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

* No cases reported for years not shown prior to 1952.

TABLE 20B.—REPORTED CASES OF SMALLPOX BY YEAR OF REPORT*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	556	17.5	326	20.7	230	14.4	177	7.9	110	9.8	67	6.0	379	40.5	216	47.5	163	34.0	
1931	63	2.0	33	2.1	30	1.9	32	1.4	15	1.3	17	1.5	31	3.3	18	3.9	13	2.7	
1932	72	2.2	38	2.3	34	2.1	45	1.9	21	1.8	24	2.1	27	2.8	17	3.7	10	2.0	
1933	35	1.1	21	1.3	14	0.8	13	0.6	6	0.5	7	0.6	22	2.3	15	3.2	7	1.4	
1934	18	0.5	14	0.8	4	0.2	10	0.4	9	0.8	1	0.1	8	0.8	5	1.1	3	0.6	
1935	24	0.7	16	1.0	8	0.5	14	0.6	8	0.7	6	0.5	10	1.0	8	1.7	2	0.4	
1936	19	0.6	13	0.8	6	0.4	5	0.2	3	0.2	2	0.2	14	1.4	10	2.1	4	0.8	
1937	11	0.3	6	0.4	5	0.3	3	0.1	1	0.1	2	0.2	8	0.8	5	1.0	3	0.6	
1938	35	1.0	22	1.3	13	0.7	17	0.7	9	0.7	8	0.6	18	1.8	13	2.7	5	1.0	
1939	15	0.4	11	0.6	4	0.2	11	0.4	7	0.6	4	0.3	4	0.4	4	0.8			
1940	6	0.2	5	0.3	1	0.1	3	0.1	2	0.2	1	0.1	3	0.3	3	0.6			
1941	4	0.1	4	0.2			3	0.1	3	0.2			1	0.1	1	0.2			
1942	7	0.2	4	0.2	3	0.2	4	0.2	1	0.1	3	0.2	3	0.3	3	0.6			
1943	21	0.6	9	0.5	12	0.6	17	0.6	6	0.5	11	0.8	4	0.4	3	0.6	1	0.2	
1944	4	0.1			4	0.2							4	0.4			4	0.8	
1945	1	0.0			1	0.1	1	0.0			1	0.1							
1946	2	0.1	1	0.1	1	0.1	1	0.0			1	0.1	1	0.1	1	0.2			
1947	2	0.1	1	0.1	1	0.1	1	0.0	1	0.1			1	0.1			1	0.2	
1948	3	0.1	3	0.2			1	0.0	1	0.1			2	0.2	2	0.4			
1951																			
1952																			

* No cases reported for years not shown prior to 1952.

TABLE 21A.—REPORTED UNTREATED CASES OF SYPHILIS BY COUNTY AND COLOR, 1951 AND 1952, WITH STAGE AND SEX, 1952
(Place of Report)

Area	1951 Reports			1952 Reports																				
							Primary-Secondary				Early latent				Late latent and other				Congenital			
							White		Other		White		Other		White		Other		White		Other	
	Total	White	Other	Total	White	Other	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F
North Carolina	4,080	459	3,621	3,709	294	3,415	33	17	193	193	52	47	546	819	82	39	722	615	9	15	144	183
Alamance	36	6	30	21	3	18				2			6	3	1	2	2	4				1
Alexander	5	1	4	5	1	4									1			1				3
Alleghany	3	1	2																			
Anson	27		27	10		10			2	3			1	2			1	1				
Ashe	1		1																			
Avery	1	1		2	2						1									1		
Beaufort	75	10	65	135	5	130				1	1	1	28	34	1		21	30	2		3	13
Bertie	33		33	25		25			1	2			7	6			3	4			2	
Bladen	37	2	35	22	1	21								5	1			7			4	5
Brunswick	15		15	19	2	17			1	1			2			1	3	6		1	4	
Buncombe	76	25	51	66	26	40	2		1	2	6	4	7	15	12	2	8	7				
Burke	64	15	49	12	6	6			2	3					1	3	1			2		
Cabarrus	45	5	40	39	5	34	1	2	7	11		1	4	7	1		1	2			1	1
Caldwell	18	6	12	10	3	7			1		1		1	2		1	2		1			1
Camden	4		4																			
Carteret	16	2	14	23		23			2					7			13	1				
Caswell	16		16	6		6			1				3								2	
Catawba	15	8	7	7	2	5								1			3		1	1	1	
Chatham	10	1	9	4	2	2						2	1				1					
Cherokee	3	3		3	3			1				1			1							

Chowan	24	2	22	12		12							3	2			3	2				2
Clay				1	1						1											
Cleveland	51	6	45	24	3	21	1	1		1			6	8	1		3	2			1	
Columbus	76	5	71	62	1	61			6	10			12	14		1	7	8			2	2
Craven	75	6	69	62	10	52	4		3	2	2	1	10	21	1	2	9	6				1
Cumberland	236	32	204	354	38	316	4		25	3	10	4	68	85	17	3	91	40			2	2
Currituck	3	1	2	10	2	8			1	1		1	1	2		1	2				1	
Dare				4		4				2							2					
Davidson	12	6	6	13	1	12			2	3	1			5			2					
Davie	10	3	7	4	2	2				1	1	1	1									
Duplin	19		19	16		16			1				1	2			1	3			4	4
Durham	108	12	96	106	7	99			16	12		1	12	12	4	2	19	19			4	5
Edgecombe	111	1	110	81	1	80			5	3			9	38			12	7		1	5	1
Forsythe	182	18	164	157	16	141	5	2	7	11		4	15	29	2		36	33	2	1	2	8
Franklin	28		28	16		16				1			3	7			1	1			1	2
Gaston	59	19	40	65	19	46	1	3	3	1	4	2	3	19	4	4	9	9		1		2
Gates	5	1	4	5		5				1				1			1	1			1	
Graham	3	3																				
Granville	30		30	15		15				1				4			6	4				
Greene	25	1	24	14	1	13				1	1		2	5			2	1			2	
Guilford	117	12	105	87	8	79	2		2	3	2	2	23	29	2		7	13				2
Halifax	60		60	166		166			5	6			22	40			33	32			10	18
Harnett	42	4	38	70	1	69			5	9			8	11	1		19	9			8	
Haywood	10	8	2	9	2	7		1	2	3						1		2				
Henderson	76	13	63	85	4	81			4	1			27	26	2	2	12	10			1	
Hertford	18	1	17	5		5			1	1				2			1					
Hoke	20		20	9		9			1	3			1	1			1	1			1	
Hyde	9		9	23		23			3	4			4	9			2				1	
Iredell	83	11	72	34	2	32		1	2	1			5	18	1		2	4				
Jackson	2	1	1	2	1	1								1	1							

TABLE 21A.—SYPHILIS—*Continued*

Area	1951 Reports			1952 Reports																							
							Primary-Secondary				Early latent				Late latent and other				Congenital			
							White		Other		White		Other		White		Other		White		Other	
	Total	White	Other	Total	White	Other	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F
Johnston	44	3	41	21	3	18	1		1	3	1	1	5	3			2	1				3
Jones	13		13	1		1								1								
Lee	27	1	26	10		10			1	1			3	2			2	1				
Lenoir	126	10	116	76	2	74			7	5			13	29		2	8	9				3
Lincoln	5	3	2	10	3	7			1					1	1	2	2	3				
McDowell	11	1	10	9	2	7			1	2		1	1		1		2	1				
Macon	7	4	3	5	3	2	1				1		1		1		1					
Madison	8	6	2	2	2						1	1										
Martin	29		29	12	1	11				1		1	2	5			1					2
Mecklenburg	202	29	173	130	13	117	3	4	1	2	1	1	12	7	1	1	37	27	1	1	12	19
Mitchell	3	3		1	1							1										
Montgomery	19	3	16	29	1	28							8	6	1		11	1			2	
Moore	36	1	35	19	1	18					1		2	7			5	3				1
Nash	82	4	78	43	1	42			1				8	16	1		5	9			1	2
New Hanover	167	13	154	227	16	211	2	1	16	4	6	3	20	35	2		68	61		2	4	3
Northampton	24	1	23	61	1	60			1				8	10	1		11	19			5	6
Onslow	107	30	77	31	8	23	5		1	1	1		12	4		1	2	1		1		2
Orange	21	2	19	10	1	9				1			1	4	1		2	1				
Pamlico	22	1	21	22		22							3	8			2	5			2	2
Pasquotank	70	3	67	166	2	164			12	10			35	38		2	31	32			1	5

Pender	30		30	42		42				2			5	6			17	9			2	1
Perquimans	19		19	15		15			1				2	1			4	5				2
Person	24		24	18		18			1				1	3			5	4			3	1
Pitt	92	5	87	47		47			8	14			2	9			2	7			1	4
Polk	3	3		2	1	1							1		1							
Randolph	16	2	14	3		3							1	1			1					
Richmond	29	1	28	29	1	28				2	1		3	8			4	5				6
Robeson	129	6	123	325	11	314			5	10	2	3	32	46	5		81	88		1	28	24
Rockingham	74	5	69	17	4	13		1	1	5		1	4	2	2			1				
Rowan	37	3	34	26	9	17	1		1	1		5	7	6	1	1	1		1			1
Rutherford	10	6	4	8	4	4					2			1	1		2	1	1			
Sampson	29	3	26	8		8				1			1	3				2			1	
Scotland	31	2	29	24	5	19			2	2		1	4	5	1	2	2	3		1		1
Stanly	20	1	19	10		10							1	2			2	5				
Stokes				1		1								1								
Surry	7	3	4	4	2	2			1							2		1				
Swain	4	2	2																			
Transylvania	4	3	1																			
Tyrrell	3	1	2	1		1											1					
Union	19	3	16	5		5			1					2				1				1
Vance	49	3	46	25		25			4	4			3	8			3	2				1
Wake	174	15	159	95	8	87			1	4	2	1	6	13	4	1	33	23			2	5
Warren	23		23	11		11								2			5	1				3
Washington	31	1	30	9		9			1				1				3	3				1
Watauga	5	5																				
Wayne	78	5	73	77	4	73			3	1	2	2	22	22			8	5			7	5
Wilkes	6	3	3	1		1															1	
Wilson	106	1	105	100	3	97			8	7			20	29	2		13	5		1	9	6
Yadkin	7	4	3	1		1											1					
Yancey	4	3	1																			

TABLE 21B.—REPORTED UNTREATED CASES OF SYPHILIS BY YEAR OF REPORT AND AGE, 1930-1952 (Place of Report)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	5,106	18	5	4	1	3	3	5	9	1	6	65	841	1,478	925	493	513	217	51	8	2	458
1931	5,144	17	4	8	4	2	4	7	5	5	6	69	843	1,595	967	596	509	189	61	10	1	242
1932	4,365	18	1	3	3	4	2	4	7	6	4	57	736	1,358	806	499	452	186	54	8	------	157
1933	4,570	21	4	3	7	1	3	7	5	11	9	61	751	1,390	854	538	516	179	79	13	3	115
1934	5,201	30	14	6	6	5	5	6	4	6	8	70	759	1,571	978	569	577	229	74	19	4	261
1935	5,189	20	9	4	7	3	7	12	6	4	10	74	774	1,556	1,030	638	575	258	95	16	3	88
1936	6,086	24	10	10	6	8	9	7	15	16	16	122	1,009	1,726	1,220	675	707	250	93	26	6	131
1937	12,299	68	14	26	25	28	32	49	45	47	54	315	1,860	3,456	2,314	1,238	1,454	454	179	47	12	582
1938	20,714	118	32	40	27	31	35	53	63	64	59	474	2,810	5,050	4,056	2,402	2,741	1,043	383	161	18	1,054
1939	30,985	115	31	59	53	43	65	90	109	65	62	629	3,631	7,303	6,727	4,223	4,874	1,760	530	189	6	421
1940	21,333	79	21	41	36	30	45	62	75	45	43	433	2,500	5,028	4,631	2,907	3,356	1,212	365	130	4	290
1941	17,560	98	19	39	28	37	38	102	38	59	29	379	2,319	4,470	3,667	2,588	2,401	769	263	91	6	120
1942	16,713	62	28	29	21	21	19	39	40	29	34	265	2,477	4,304	3,065	2,347	2,787	793	238	60	5	50
1943	14,125	32	44	12	13	20	12	20	32	30	29	242	2,910	3,564	2,392	1,926	1,994	561	184	57	6	45
1944	9,804	46	24	21	13	15	11	16	18	20	17	227	2,299	2,570	1,535	1,143	1,152	454	155	62	6	--------
1945	8,125	46	77	20	10	13	9	19	15	20	25	233	2,090	1,938	1,254	755	950	430	164	50	5	2
1946	9,233	*	84	19	12	11	6	21	17	14	21	250	2,234	2,871	1,491	767	825	373	123	56	8	30
1947	8,724	103	12	12	19	6	10	14	15	9	16	249	2,076	2,660	1,422	643	761	416	159	72	18	32
1948	7,313	91	15	12	12	10	11	14	10	11	11	177	1,618	2,220	1,170	632	730	333	147	61	6	22
1949	6,699	65	30	10	7	8	10	15	14	12	20	205	1,403	1,750	1,046	634	781	438	156	56	16	23
1950	5,146	50	7	9	13	8	7	9	7	9	16	156	953	1,273	848	522	653	352	155	67	11	21
1951	4,080	33	3	9	13	7	13	12	12	9	9	188	768	874	564	396	537	357	177	70	11	18
1952	3,709	23	5	3	3	3	3	5	8	18	11	150	568	538	446	381	694	458	247	108	19	18

* "Under 1 yr." combined with "1 yr." group.

TABLE 21C.—REPORTED UNTREATED CASES OF SYPHILIS BY YEAR OF REPORT, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952 (Place of Report)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	5,106	161.0	2,705	171.7	2,401	150.5	922	41.2	602	53.7	370	33.2	4,134	442.1	2,103	462.3	2,031	423.1	----------
1931	5,144	160.4	2,816	176.7	2,327	144.3	969	42.7	567	49.9	402	35.6	4,170	443.7	2,245	490.8	1,925	399.1	5
1932	4,365	133.7	2,222	136.9	2,143	130.5	912	39.4	551	47.5	361	31.3	3,438	361.5	1,661	358.7	1,777	364.3	15
1933	4,570	137.7	2,461	149.2	2,096	125.6	805	34.1	462	39.1	343	29.2	3,759	391.2	1,993	425.6	1,753	355.9	13
1934	5,201	154.8	2,517	150.7	2,684	158.9	992	41.5	594	49.6	398	33.4	4,203	434.5	1,920	406.9	2,283	460.7	6
1935	5,189	153.5	2,491	148.2	2,698	158.8	878	36.4	544	45.0	334	27.8	4,307	443.9	1,943	410.3	2,364	475.9	4
1936	6,086	179.0	2,645	156.5	3,441	201.2	951	39.2	536	44.1	415	34.2	5,120	526.1	2,098	441.4	3,022	607.0	15
1937	12,299	358.4	4,911	287.9	7,388	427.9	1,735	70.7	785	63.9	950	77.5	10,501	1,073.4	4,089	855.1	6,412	1,282.2	63
1938	20,714	596.0	8,143	471.4	12,571	719.1	2,698	108.3	1,321	106.0	1,377	110.6	17,736	1,801.0	6,756	1,402.3	10,980	2,183.0	280
1939	30,985	877.3	13,757	783.7	17,228	969.8	4,431	174.7	2,386	188.2	2,045	161.2	26,554	2,668.6	11,371	2,333.4	15,183	1,990.2	----------
1940	21,333	597.3	9,365	528.2	11,968	665.4	2,764	107.6	1,433	111.8	1,331	103.5	18,569	1,849.5	7,932	1,614.7	10,637	2,074.5	----------
1941	17,560	478.5	8,864	486.9	8,696	470.1	2,143	81.3	1,198	91.2	945	71.5	15,417	1,489.6	7,666	1,513.6	7,751	1,466.7	----------
1942	16,713	456.0	9,996	554.3	6,717	360.7	2,000	76.0	1,302	100.1	698	52.5	14,713	1,423.4	8,694	1,731.3	6,019	1,132.5	----------
1943	14,125	383.5	7,835	437.7	6,290	332.3	1,814	68.8	998	77.9	816	60.2	12,311	1,177.0	6,837	1,345.2	5,474	1,018.0	----------
1944	9,804	272.7	4,533	266.0	5,271	278.8	1,550	60.2	777	64.1	773	56.7	8,254	808.5	3,756	762.6	4,498	851.2	----------
1945	8,125	231.2	3,618	220.9	4,507	240.2	1,536	61.1	795	68.8	741	54.6	6,589	657.9	2,823	586.3	3,766	724.1	----------
1946	9,233	253.6	4,369	245.4	4,864	261.5	1,615	62.0	950	74.3	665	50.0	7,618	736.8	3,419	680.5	4,199	790.0	----------
1947	8,724	234.6	3,897	213.5	4,827	255.1	1,438	54.0	793	60.4	645	47.8	7,286	690.0	3,104	605.0	4,182	770.4	----------
1948	7,313	192.5	3,308	177.4	4,005	207.2	1,220	44.9	684	51.0	536	38.9	6,093	564.9	2,624	500.6	3,469	625.6	----------
1949	6,699	173.4	2,981	157.1	3,718	189.0	1,016	36.7	570	41.8	446	31.8	5,683	517.9	2,411	452.1	3,272	580.0	----------
1950	5,146	126.7	2,258	111.9	2,888	141.2	498	16.7	252	16.9	246	16.5	4,648	430.8	2,006	380.4	2,642	479.0	----------
1951	4,080	99.0	1,871	91.4	2,209	106.4	459	15.1	251	16.6	208	13.7	3,621	332.8	1,620	304.6	2,001	359.6	----------
1952	3,709	88.9	1,781	86.0	1,928	91.8	294	9.6	176	11.5	118	7.7	3,415	311.7	1,605	299.8	1,810	323.1	----------

TABLE 21D.—REPORTED UNTREATED CASES OF SYPHILIS BY COLOR, STAGE, AND SEX WITH RATES PER 100,000 POPULATION, 1940-1952*

Year	Color	All Stages			Primary-Secondary			Early Latent			Late and Unknown			Congenital		
		Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
	Total	21,225	9,528	11,697	2,611	1,307	1,304	9,436	3,766	5,670	8,143	4,009	4,134	1,035	446	589
	Rate	594.3	537.4	650.3	73.1	73.7	72.5	264.2	212.4	315.2	228.0	226.1	229.8	29.0	25.2	32.7
1940	White	2,756	1,454	1,302	504	280	224	986	467	519	1,104	632	472	162	75	87
	Rate	107.3	113.4	101.3	19.6	21.8	17.4	38.4	36.4	40.4	43.0	49.3	36.7	6.3	5.9	6.8
	Other	18,469	8,074	10,395	2,107	1,027	1,080	8,450	3,299	5,151	7,039	3,377	3,662	873	371	502
	Rate	1,839.6	1,643.6	2,027.3	209.9	209.1	210.6	841.6	671.6	1,004.6	701.1	687.5	714.2	87.0	75.5	97.9
	Total	18,627	9,403	9,224	3,805	2,087	1,718	7,357	3,515	3,842	6,567	3,421	3,146	898	380	518
	Rate	507.5	516.6	498.7	103.7	114.6	92.9	200.5	193.1	207.7	178.9	187.9	170.1	24.5	20.9	28.0
1941	White	2,270	1,272	998	445	247	198	705	386	319	971	568	403	149	71	78
	Rate	86.1	96.8	75.5	16.9	18.8	15.0	26.8	29.4	24.1	36.8	43.2	30.5	5.7	5.4	5.9
	Other	16,357	8,131	8,226	3,360	1,840	1,520	6,652	3,129	3,523	5,596	2,853	2,743	749	309	440
	Rate	1,580.5	1,605.5	1,556.5	324.7	363.3	287.6	642.7	617.8	666.6	540.7	563.3	519.0	72.4	61.0	83.3
	Total	16,695	9,986	6,709	3,616	2,158	1,458	6,819	3,743	3,076	5,799	3,876	1,923	461	209	252
	Rate	455.5	553.7	360.3	98.7	119.7	78.3	186.0	207.6	165.2	158.2	214.9	103.3	12.6	11.6	13.5
1942	White	2,004	1,304	700	518	343	175	607	352	255	802	570	232	77	39	38
	Rate	76.1	100.2	52.6	19.7	26.4	13.2	23.1	27.1	19.2	30.5	43.8	17.4	2.9	3.0	2.9
	Other	14,691	8,682	6,009	3,098	1,815	1,283	6,212	3,391	2,821	4,997	3,306	1,691	384	170	214
	Rate	1,421.3	1,728.9	1,130.6	299.7	361.4	241.4	601.0	675.3	530.8	483.4	658.4	318.2	37.2	33.9	40.3

	Total	13,709	7,627	6,082	3,267	1,755	1,512	5,866	2,969	2,897	4,324	2,787	1,537	252	116	136
	Rate	372.2	426.1	321.3	88.7	98.0	79.9	159.3	165.9	153.0	117.4	155.7	81.2	6.8	6.5	7.2
1943	White	1,628	904	724	477	276	201	536	258	278	568	348	220	47	22	25
	Rate	61.7	70.5	53.4	18.1	21.5	14.8	20.3	20.1	20.5	21.5	27.2	16.2	1.8	1.7	1.8
	Other	12,081	6,723	5,358	2,790	1,479	1,311	5,330	2,711	2,619	3,756	2,439	1,317	205	94	111
	Rate	1,155.0	1,322.8	996.4	266.7	291.0	243.8	509.6	533.4	487.1	359.1	479.9	244.9	19.6	18.5	20.6
	Total	9,823	4,542	5,281	3,350	1,721	1,629	3,767	1,353	2,414	2,528	1,389	1,139	178	79	99
	Rate	273.3	266.6	279.3	93.2	101.0	86.1	104.8	79.4	127.7	70.3	81.5	60.2	5.0	4.6	5.2
1944	White	1,556	780	776	734	406	328	426	144	282	356	212	144	40	18	22
	Rate	60.5	64.4	57.0	28.5	33.5	24.1	16.6	11.9	20.7	13.8	17.5	10.6	1.6	1.5	1.6
	Other	8,267	3,762	4,505	2,616	1,315	1,301	3,341	1,209	2,132	2,172	1,177	995	138	61	77
	Rate	809.7	763.9	852.5	256.2	267.0	246.2	327.2	245.5	403.4	212.7	239.0	188.3	13.5	12.4	14.6
	Total	8,212	3,664	4,548	3,037	1,508	1,529	3,072	1,093	1,979	1,691	876	815	412	187	225
	Rate	233.7	223.7	242.3	86.4	92.1	81.5	87.4	66.7	105.5	48.1	53.5	43.4	11.7	11.4	12.0
1945	White	1,570	810	760	765	451	314	415	149	266	323	180	143	67	30	37
	Rate	62.5	70.1	56.0	30.4	39.0	23.1	16.5	12.9	19.6	12.9	15.6	10.5	2.7	2.6	2.7
	Other	6,642	2,854	3,788	2,272	1,057	1,215	2,657	944	1,713	1,368	696	672	345	157	188
	Rate	663.2	592.8	728.3	226.8	219.5	233.6	265.3	196.1	329.3	136.6	144.6	129.2	34.4	32.6	36.1
	Total	9,262	4,387	4,875	4,574	2,452	2,122	3,101	1,153	1,948	1,155	574	581	432	208	224
	Rate	254.4	246.4	262.0	125.6	137.7	114.1	85.2	64.8	104.7	31.7	32.2	31.2	11.9	11.7	12.0
1946	White	1,620	954	666	1,018	660	358	337	143	194	201	121	80	64	30	34
	Rate	62.1	74.7	50.1	39.1	51.6	26.9	12.9	11.2	14.6	7.7	9.5	6.0	2.5	2.3	2.6
	Other	7,642	3,433	4,209	3,556	1,792	1,764	2,764	1,010	1,754	954	453	501	368	178	190
	Rate	739.1	683.3	791.9	343.9	356.7	331.9	267.3	201.0	330.0	92.3	90.2	94.3	35.6	35.4	35.7

TABLE 21D.—SYPHILIS—*Continued*

Year	Color	All Stages			Primary-Secondary			Early Latent			Late and Unknown			Congenital		
		Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
	Total	8,760	3,912	4,848	4,072	2,082	1,990	3,205	1,080	2,125	1,112	578	534	371	172	199
	Rate	235.6	214.3	256.2	109.5	114.0	105.2	86.2	59.2	112.3	29.9	31.7	28.2	10.0	9.4	10.5
1947	White	1,440	793	647	859	514	345	336	129	207	184	119	65	61	31	30
	Rate	54.1	60.4	47.9	32.3	39.2	25.6	12.6	9.8	15.3	6.9	9.1	4.8	2.3	2.4	2.2
	Other	7,320	3,119	4,201	3,213	1,568	1,645	2,869	951	1,918	928	459	469	310	141	169
	Rate	693.2	607.9	773.9	304.3	305.6	303.0	271.7	185.3	353.3	87.9	89.5	86.4	29.4	27.5	31.1
	Total	7,313	3,308	4,005	2,992	1,627	1,365	2,905	1,025	1,880	1,060	502	558	356	154	202
	Rate	192.5	177.4	207.2	78.8	87.2	70.6	76.5	55.0	97.2	27.9	26.9	28.9	9.4	8.3	10.4
1948	White	1,220	684	536	673	433	240	315	124	191	186	106	80	46	21	25
	Rate	44.9	51.0	38.9	24.7	32.3	17.4	11.6	9.2	13.9	6.8	7.9	5.8	1.7	1.6	1.8
	Other	6,093	2,624	3,469	2,319	1,194	1,125	2,590	901	1,689	874	396	478	310	133	177
	Rate	564.9	500.6	625.6	215.0	227.8	202.9	240.1	171.9	304.6	81.0	75.6	86.2	28.7	25.4	31.9
	Total	6,699	2,981	3,718	2,093	1,115	978	2,935	1,062	1,873	1,254	617	637	417	187	230
	Rate	173.4	157.1	189.0	54.2	58.8	49.7	76.0	56.0	95.2	32.5	32.5	32.4	10.8	9.9	11.7
1949	White	1,016	570	446	409	277	132	316	126	190	248	146	102	43	21	22
	Rate	36.7	41.8	31.8	14.8	20.3	9.4	11.4	9.2	13.5	9.0	10.7	7.3	1.6	1.5	1.6
	Other	5,683	2,411	3,272	1,684	838	846	2,619	936	1,683	1,006	471	535	374	166	208
	Rate	517.9	452.1	580.0	153.5	157.1	150.0	238.7	175.5	298.4	91.7	88.3	94.8	34.1	31.1	36.9
	Total	5,146	2,258	2,888	1,187	590	597	2,457	922	1,535	1,170	595	575	332	151	181
	Rate	126.7	111.9	141.2	29.2	29.2	29.2	60.5	45.7	75.1	28.8	29.5	28.1	8.2	7.5	8.9
1950	White	498	252	246	163	99	64	168	68	100	139	74	65	28	11	17
	Rate	16.7	16.9	16.5	5.5	6.6	4.3	5.6	4.6	6.7	4.7	5.0	4.4	0.9	0.7	1.1
	Other	4,648	2,006	2,642	1,024	491	533	2,289	854	1,435	1,031	521	510	304	140	164
	Rate	430.8	380.5	479.0	94.9	93.1	96.6	212.2	162.0	260.2	95.6	98.8	92.5	28.2	26.6	29.7

	Total	4,080	1,872	2,208	744	407	337	1,804	707	1,097	1,114	570	544	418	188	230
	Rate	99.0	91.4	106.4	18.0	19.9	16.2	43.8	34.5	52.9	27.0	27.8	26.2	10.1	9.2	11.1
	White	459	251	208	118	78	40	122	57	65	172	95	77	47	21	26
1951	Rate	15.1	16.6	13.7	3.9	5.1	2.6	4.0	3.8	4.3	5.7	6.3	5.1	1.5	1.4	1.7
	Other	3,621	1,621	2,000	626	329	297	1,682	650	1,032	942	475	467	371	167	204
	Rate	332.8	304.8	359.5	57.5	61.9	53.4	154.6	122.2	185.5	86.6	89.3	83.9	34.1	31.4	36.7
	Total	3,709	1,781	1,928	436	226	210	1,464	598	866	1,458	804	654	351	153	198
	Rate	88.9	86.0	91.8	10.4	10.9	10.0	35.1	28.9	41.2	34.9	38.8	31.1	8.4	7.4	9.4
	White	294	176	118	50	33	17	99	52	47	121	82	39	24	9	15
1952	Rate	9.6	11.5	7.7	1.6	2.1	1.1	3.2	3.4	3.1	3.9	5.3	2.5	0.8	0.6	1.0
	Other	3,415	1,605	1,810	386	193	193	1,365	546	819	1,337	722	615	327	144	183
	Rate	311.7	299.8	323.1	35.2	36.0	34.4	124.6	102.0	146.2	122.0	134.9	109.8	29.8	26.9	32.7

* Source: Monthly venereal disease morbidity reports, USPHS Report Form 8958-B, 1940-1947. Cases recorded by place of report rather than by place of residence. The totals for these years will not necessarily agree with other similar totals in this bulletin or in previous bulletins because the data have been corrected monthly; whereas, other data have been corrected monthly and annually.

TABLE 21E.—REPORTED CASES OF CONGENITAL SYPHILIS, AND YEAR OF BIRTH: NORTH

Year of Birth	Year of Report											
	Total	White	Other	1952			1951			1950		
				Total	White	Other	Total	White	Other	Total	White	Other
1952	22	------	22	22	------	22	------	------	------	------	------	------
1951	37	9	28	5	------	5	32	9	23	------	------	------
1950	56	3	53	3	1	2	3	------	3	50	2	48
1949	80	13	67	3	------	3	9	2	7	6	------	6
1948	130	13	117	2	------	2	11	2	9	8	1	7
1947	138	15	123	3	------	3	7	------	7	12	------	12
1946	124	20	104	5	------	5	13	2	11	6	1	5
1945	124	22	102	8	1	7	11	1	10	6	1	5
1944	94	6	88	17	1	16	12	------	12	9	------	9
1943	81	13	68	10	3	7	9	1	8	7	------	7
1942	78	5	73	13	------	13	9	1	8	8	1	7
1941	89	11	78	14	------	14	18	1	17	13	4	9
1940	123	14	109	27	2	25	20	3	17	14	1	13
1939	129	13	116	26	2	24	24	4	20	19	------	19
1938	151	14	137	35	1	34	40	2	38	13	1	12
1937	187	22	165	36	1	35	31	------	31	19	3	16
1936	200	24	176	42	3	39	39	4	35	17	------	17
1935	188	24	164	25	4	21	22	4	18	31	1	30
1934	190	18	172	15	1	14	32	3	29	21	3	18
1933	163	21	142	10	------	10	18	2	16	23	4	19
1932	155	18	137	4	------	4	20	1	19	17	1	16
1931	141	19	122	6	1	5	13	2	11	10	------	10
1930	94	9	85	3	------	3	7	2	5	4	------	4
1929	70	4	66	3	------	3	1	------	1	3	------	3
1928	45	6	39	1	------	1	5	------	5	4	1	3
1927	38	8	30	3	------	3	2	------	2	1	------	1
1926	25	4	21	1	------	1	1	------	1	1	------	1
1925	16	1	15	1	------	1	2	------	2	1	------	1
Prior to 1925	113	26	87	7	3	4	6	------	6	9	3	6
Total	3,081	375	2,706	350	24	326	417	46	371	332	28	304

PREVIOUSLY UNTREATED, BY COLOR SHOWING YEAR OF REPORT, 1945-1952
CAROLINA, ALL REPORT SOURCES

Year of Report														
1949			1948			1947			1946			1945		
Total	White	Other	Total	White	Other	Total	White	Other	Total	White	Other	Total	White	Other
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------
62	11	51	------	------	------	------	------	------	------	------	------	------	------	------
27	2	25	82	8	74	------	------	------	------	------	------	------	------	------
10	------	10	20	1	19	86	14	72	------	------	------	------	------	------
5	1	4	8	------	8	21	1	20	66	15	51	------	------	------
7	1	6	10	1	9	5	------	5	17	1	16	60	16	44
10	2	8	6	1	5	11	1	10	14	------	14	15	1	14
11	1	10	9	2	7	6	------	6	11	2	9	18	4	14
11	------	11	13	2	11	6	------	6	10	1	9	8	------	8
11	2	9	9	3	6	8	------	8	6	------	6	10	1	9
17	2	15	8	------	8	12	2	10	17	2	15	8	2	6
11	1	10	9	1	8	6	1	5	17	3	14	17	1	16
13	2	11	15	4	11	12	2	10	10	------	10	13	2	11
28	3	25	16	4	12	21	5	16	21	3	18	15	3	12
21	2	19	10	2	8	25	5	20	27	4	23	19	4	15
32	5	27	20	2	18	14	5	9	23	1	22	21	2	19
32	------	32	23	1	22	24	6	18	28	1	27	15	3	12
20	2	18	25	3	22	15	3	12	22	1	21	30	6	24
20	2	18	17	2	15	21	4	17	26	6	20	30	2	28
24	------	24	12	3	9	23	4	19	30	8	22	23	1	22
16	------	16	11	1	10	13	------	13	14	3	11	26	3	23
3	1	2	8	------	8	16	1	15	16	1	15	20	1	19
1	------	1	3	------	3	6	2	4	15	1	14	10	2	8
3	1	2	1	------	1	2	------	2	11	3	8	15	4	11
5	------	5	2	------	2	3	1	2	7	3	4	5	------	5
6	1	5	3	------	3	------	------	------	2	------	2	1	------	1
11	1	10	16	5	11	14	3	11	20	5	15	30	6	24
417	43	374	356	46	310	370	60	310	430	64	366	409	64	345

TABLE 22A.—REPORTED CASES OF TETANUS BY COUNTY AND COLOR WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1952 Onsets																
	Color				Month of onset												
	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	26	8	18			4	2	2	1	1	7	4	1	1		1	2
Beaufort	2	1	1								2						
Buncombe	1	1										1					
Caldwell	1	1									1						
Cleveland	1	1								1							
Durham	1		1								1						
Edgecombe	3		3									3					
Johnston	1	1				1											
Lenoir	1		1								1						
Martin	2	1	1			1					1						
Mecklenburg	1	1				1											
Nash	1		1					1									
New Hanover	3		3			1		1	1								
Pasquotank	2	1	1											1			1
Pitt	5		5				1				1		1			1	1
Richmond	1		1				1										

TABLE 22B.—REPORTED CASES OF TETANUS BY YEAR AND AGE, 1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and Over	Not stated
1952	26	7	------	1	3	3	1	2	------	------	------	3	1	------	------	------	2	2	------	1	------	------

TABLE 22C.—REPORTED CASES OF TETANUS BY YEAR, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1952 (Place of Residence)

Year	All Classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1952	26	0.6	20	1.0	6	0.3	8	0.3	8	0.5	------	------	18	1.6	12	2.2	6	1.1	------

TABLE 23A.—REPORTED CASES OF TRACHOMA BY YEAR OF REPORT* AND AGE, 1931-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1931	6		1		1							3				1						
1933	5	1	1		2									1								
1934	1																		1			
1935	6					1	2			1	1						1					
1936	1									1												
1940	8					1		1	1			2		1					1	1		
1941	4			1		1					1	1										
1943	1													1								
1945	1								1													
1952																						

* No cases reported for years not shown prior to 1952.

Table 23B.—Reported Cases of Trachoma by Year of Report*, Color, and Sex with Rates per 100,000 Population, 1931-1952 (Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1931	6	0.2	3	0.2	3	0.2	6	0.3	3	0.3	3	0.3	------	------	------	------	------	------	------
1933	5	0.2	3	0.2	2	0.1	5	0.2	3	0.3	2	0.2	------	------	------	------	------	------	------
1934	1	0.0	1	0.1	------	------	1	0.0	1	0.1	------	------	------	------	------	------	------	------	------
1935	6	0.2	3	0.2	3	0.2	6	0.2	3	0.2	3	0.2	------	------	------	------	------	------	------
1936	1	0.0	1	0.1	------	------	1	0.0	1	0.1	------	------	------	------	------	------	------	------	------
1940	8	0.2	4	0.2	4	0.2	4	0.2	2	0.2	2	0.2	4	0.4	2	0.4	2	0.4	------
1941	4	0.1	3	0.2	1	0.1	2	0.1	1	0.1	1	0.1	2	0.2	2	0.4	------	------	------
1943	1	0.0	1	0.1	------	------	1	0.0	1	0.1	------	------	------	------	------	------	------	------	------
1945	1	0.0	1	0.1	------	------	------	------	------	------	------	------	1	0.1	1	0.2	------	------	------
1952	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

* No cases reported for years not shown prior to 1952.

TABLE 24A.—REPORTED CASES OF TUBERCULOSIS, ALL FORMS, BY COUNTY, (Place of

Area	1951 Reports											
	Respiratory			Other forms			Other forms		Total		Childhood	
											Inflammatory	
	Total	White	Other	Total	White	Other	White	Other	White	Other	White	Other
North Carolina	2,940	1,801	1,139	165	69	96	66	69	1,280	911	67	94
Alamance	35	21	14	5	3	2	2	------	8	9	------	------
Alexander	10	10	------	------	------	------	------	------	2	1	------	------
Alleghany	5	5	------	1	1	------	------	------	5	1	1	------
Anson	8	1	7	------	------	------	------	2	2	5	------	2
Ashe	21	21	------	1	1	------	2	------	9	------	------	------
Avery	11	11	------	1	1	------	------	------	10	------	1	------
Beaufort	24	12	12	3	1	2	1	2	7	14	------	1
Bertie	14	7	7	5	4	1	------	------	15	11	1	2
Bladen	13	7	6	------	------	------	------	1	5	6	------	2
Brunswick	10	4	6	2	1	1	------	1	2	1	------	------
Buncombe	254	189	65	12	8	4	3	------	86	33	2	1
Burke	39	30	9	2	2	------	1	1	12	2	2	------
Cabarrus	28	23	5	2	1	1	1	1	44	6	2	------
Caldwell	14	12	2	------	------	------	2	1	32	5	------	1
Camden	4	4	------	2	1	1	------	------	------	------	------	------
Carteret	15	5	10	------	------	------	------	------	7	4	------	2
Caswell	8	7	1	------	------	------	------	1	5	4	1	1
Catawba	12	11	1	2	2	------	------	------	16	1	------	------
Chatham	6	3	3	------	------	------	------	------	1	3	------	1
Cherokee	4	4	------	1	1	------	------	------	16	1	1	------
Chowan	10	1	9	------	------	------	------	------	4	5	1	------
Clay	4	4	------	------	------	------	------	------	3	------	1	------
Cleveland	30	20	10	2	1	1	1	------	9	9	------	------
Columbus	54	39	15	1	------	1	------	1	9	9	------	2
Craven	34	16	18	3	1	2	------	------	6	13	2	1
Cumberland	43	26	17	6	3	3	2	------	29	20	3	------
Currituck	5	4	1	------	------	------	------	------	1	------	------	------
Dare	4	3	1	------	------	------	------	------	2	------	------	------
Davidson	19	12	7	1	1	------	1	------	8	6	------	------
Davie	2	2	------	------	------	------	------	------	5	------	------	------
Duplin	19	10	9	1	------	1	------	1	11	5	1	2
Durham	107	67	40	10	3	7	1	2	26	39	------	2
Edgecombe	65	33	32	------	------	------	------	2	6	32	------	6
Forsyth	147	69	78	2	1	1	4	3	74	42	2	1
Franklin	67	43	24	------	------	------	------	1	7	8	------	1

COLOR, AND TYPE, 1951 AND 1952, WITH STAGE AND ACTIVITY, 1952
Residence)

1952 Reports

Respiratory

Childhood		Active						Inactive						Not stated	
Healed		Minimal		Moderate		Far advanced		Minimal		Moderate		Far advanced			
White	Other	White	Other	White	Other	White	Other	White	Other	White	Other	White	Other	White	Other
80	30	126	60	336	239	289	380	182	34	109	22	35	15	56	37
				2	1	3	5	1	1	2					2
						1	1							1	
		1		1	1					2					
				1	2	1									1
4		1		1		1		1						1	
1						1		4		2				1	
		2		1	3	3	8						1	1	1
	1			1	3	1	2	4		6	1	1	1	1	1
		2		2	1	1	3								
				1		1	1								
1		7	4	17	7	31	13	5	1	13	2	7	1	3	4
		2		3		2	2	1		1		1			
4	1	1	1	10		3	3	9		8		1		6	1
		3		4	2	2		16	2	6				1	
3		1		2			2	1							
			2	2		2	1								
		2		3		4		3		1	1	1		2	
					1	1	1								
		2		5	1	2		3		2				1	
				2	2		2	1	1						
				1		1									
		1	1	4	3	1	3	1	1			1	1	1	
		1	1	2	3	4	3			1		1			
			1	1	2	3	8		1						
		4		14	12	8	7								1
						1									
						1								1	
				2	2	2	2	3					1	1	1
		1		2		2									
		3		4		2	3	1							
		3	7	10	14	2	15	4		5	1	1		1	
	1	1		3	11		11	2	1		1				1
15	8	4	3	4	7	8	13	27	9	13	1			1	
		1		1	2	3	4		1	1		1			

TABLE 24A.—

Area	1951 Reports											
	Respiratory			Other forms			Other forms		Total		Childhood	
											Inflammatory	
	Total	White	Other	Total	White	Other	White	Other	White	Other	White	Other
Gaston	97	81	16	1	------	1	------	1	51	12	5	3
Gates	3	1	2	1	------	1	------	------	------	1	------	------
Graham	4	4	------	------	------	------	------	------	7	2	1	1
Granville	30	16	14	2	------	2	1	------	20	22	1	4
Greene	18	5	13	1	------	1	------	------	2	3	------	------
Guilford	64	29	35	3	------	3	5	6	58	33	5	------
Halifax	100	47	53	------	------	------	2	1	7	25	------	1
Harnett	43	22	21	3	1	2	1	------	17	15	3	2
Haywood	9	7	2	2	2	------	1	------	13	------	2	------
Henderson	34	32	2	2	2	------	1	------	24	1	------	------
Hertford	12	6	6	1	------	1	------	------	4	13	------	1
Hoke	10	2	8	1	------	1	------	2	4	4	1	1
Hyde	8	6	2	------	------	------	1	1	2	------	1	------
Iredell	20	19	1	------	------	------	------	------	15	10	------	3
Jackson	2	2	------	1	1	------	1	------	6	------	2	------
Johnston	31	16	15	1	1	------	1	1	36	7	1	------
Jones	6	4	2	1	------	1	------	1	1	5	------	------
Lee	21	10	11	2	1	1	------	------	6	8	------	------
Lenoir	31	13	18	4	2	2	1	------	18	23	1	1
Lincoln	5	3	2	1	------	1	------	------	3	2	------	1
McDowell	7	7	------	------	------	------	------	------	7	------	------	------
Macon	8	8	------	------	------	------	1	------	6	------	------	------
Madison	16	16	------	------	------	------	------	------	14	------	------	------
Martin	20	13	7	------	------	------	------	------	11	29	------	1
Mecklenburg	125	71	54	6	1	5	4	1	33	29	------	2
Mitchell	51	51	------	1	1	------	2	------	7	------	------	------
Montgomery	3	2	1	------	------	------	------	1	1	2	------	------
Moore	7	2	5	------	------	------	------	1	6	5	------	1
Nash	43	16	27	3	------	3	------	3	18	33	------	8
New Hanover	74	44	30	6	1	5	1	2	22	29	1	2
Northampton	21	8	13	3	1	2	1	2	5	13	1	1
Onslow	18	16	2	1	------	1	1	1	19	5	1	------
Orange	24	12	12	1	------	1	1	------	4	2	------	------
Pamlico	37	20	17	------	------	------	------	------	1	2	------	------
Pasquotank	26	15	11	3	------	3	------	------	4	7	------	------

TUBERCULOSIS—*Continued*

1952 Reports

Respiratory

Childhood		Active						Inactive						Not stated	
Healed		Minimal		Moderate		Far advanced		Minimal		Moderate		Far advanced			
White	Other	White	Other	White	Other	White	Other	White	Other	White	Other	White	Other	White	Other
6	1	7	1	13	3	6	4	10		3				1	
							1								
		1		2	1	2								1	
1	2	1	2	3	1	2	1	8	6	1		1	2	2	4
				1		1	3								
4	1	5	3	16	5	14	20	8	1	4	2	1	1	1	
		2	1	2	11	2	11	1					1		
		1		6	2	4	10					1		2	1
1				3		6								1	
1		1		6	1	5		2		8		1			
		2	1	1	6		4			1			1		
			1	1		2	2								
		1													
		4		2	2	6	5	2		1					
		1				1		1		1					
1		1		16	1	12	6	1		3		1			
				1	2		3								
				2	3	3	4	1			1				
	2	1		9	7	7	9						1		3
	1	1				1		1							
				2		2		2						1	
				2		1		1				1		1	
		1		6		5		1		1					
	1		2	3	10	5	11		1	2	3			1	
		7	1	9	7	13	16	4			1				2
		1		2		2		1				1			
						1	2								
		1	1	2	1	1	1	1	1	1					
		2	1	7	8	2	14	3		2	1			2	1
2	1	4	6	9	6	1	12	2	1	1	1	2			
				1	7	1	5			1				1	
2		3	1	3		5	4	3		2					
	1	1		1	1					1		1			
							2					1			
		1	3	1	3	2	1								

TABLE 24A.—

Area	1951 Reports											
	Respiratory			Other forms			Other forms		Total		Childhood	
											Inflammatory	
	Total	White	Other	Total	White	Other	White	Other	White	Other	White	Other
Pender	8	1	7	2	2				2	7		
Perquimans	8	4	4	1		1			6	2		
Person	23	11	12						8	9	3	5
Pitt	53	19	34	6	1	5	1	3	9	26		3
Polk	6	5	1						6		1	
Randolph	22	18	4				1		22	2	1	
Richmond	12	6	6	2	1	1		1	6	6		1
Robeson	52	20	32	14	5	9	2	3	16	26		5
Rockingham	38	24	14				1		15	18		3
Rowan	22	15	7	1		1			20	9		
Rutherford	22	19	3						8	1	1	
Sampson	38	21	17	3	1	2	1	2	12	13	1	1
Scotland	24	16	8	1		1			6	14		2
Stanly	22	19	3						10	3		2
Stokes	3	3							5	2		
Surry	36	35	1	2	2		4	1	70	2	1	
Swain	7	5	2	2		2			5	1	1	
Transylvania	6	6		1	1				7	1	1	
Tyrrell	5	3	2	1		1			2			
Union	13	8	5				2	2	6	9	1	1
Vance	21	11	10						7	9	2	1
Wake	69	34	35	8	2	6	4	6	28	26	3	1
Warren	13	5	8					1	1	5		
Washington	4	3	1					1	2	3		
Watauga	12	12					1		12		1	
Wayne	38	12	26	2	1	1		2	14	23	1	
Wilkes	23	18	5				1	1	10	5	1	
Wilson	150	70	80	2	1	1	1	2	20	51		8
Yadkin	8	7	1						2	1		
Yancey	9	9		1	1				7		1	

TUBERCULOSIS—*Continued*

1952 Reports

Respiratory

Childhood		Active						Inactive						Not stated	
Healed		Minimal		Moderate		Far advanced		Minimal		Moderate		Far advanced			
White	Other	White	Other	White	Other	White	Other	White	Other	White	Other	White	Other	White	Other
		1			2		4				1			1	
4	1	1		1	1										
					1	4	3					1			
1		1	1	3	5	4	15						1		1
						1		1						3	
2		1	1	5		4		6		2				1	1
		2		3		1	5								
				13	3	3	16		1						1
		2		3	2	3	10	4		1	1			2	2
		3	2	6	1	3	3	2		2	1	3	2	1	
2				2		3	1								
		2		4	6	3	5							2	1
		2	3	1	3	1	5	1		1			1		
				3		6	1							1	
1		1	1		1	2								1	
21		5		15	2	8		17		2				1	
				3		1	1								
		2		2	1							1		1	
				1										1	
			1	1	2	2	4	2							1
				2	2	2	6							1	
		2	1	13	10	7	9		3	2		1			2
			2		1		2	1							
		1		1	1		2								
3						1		2				2		3	
		4	2	4	13	2	4	1		1		1			4
		1		1	4	5	1	2							
	8	1	2	9	10	6	19	2	1	2	3				
		1		1					1						
				1		4		1							

TABLE 24B.—REPORTED CASES OF TUBERCULOSIS, ALL FORMS, BY YEAR OF REPORT AND AGE, 1933-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1933	2,524	------	5	8	5	9	14	26	38	44	48	268	300	384	354	226	359	183	78	32	4	139
1934	2,047	------	3	9	1	8	17	21	24	57	52	336	228	271	229	161	227	119	64	22	5	193
1935	1,996	6	4	3	9	9	17	27	37	48	50	270	191	288	277	175	256	120	62	21	7	119
1936	3,008	10	8	7	10	15	14	35	31	35	44	297	312	423	394	307	473	231	146	59	21	136
1937	2,632	12	6	4	8	9	9	5	7	2	10	78	236	426	455	270	415	269	154	73	12	172
1938	2,607	16	17	15	17	11	7	13	14	12	14	97	250	374	415	310	395	245	130	72	15	168
1939	2,366	11	12	12	9	11	16	21	24	29	24	244	291	307	315	242	340	198	129	46	4	81
1940	2,084	9	9	10	5	6	6	10	12	15	16	95	198	319	322	236	333	243	112	52	8	68
1941	2,425	7	6	13	5	12	13	12	15	20	18	112	238	335	325	271	397	260	141	62	3	160
1942	1,902	8	3	7	12	10	4	11	9	8	5	66	155	334	265	211	310	227	103	47	8	99
1943	1,633	10	8	10	7	4	4	12	6	6	10	53	178	236	218	197	258	179	103	48	8	78
1944	1,804	5	5	9	14	6	10	7	9	9	6	68	131	256	246	246	287	235	146	58	19	32
1945	3,392	4	15	22	16	33	39	69	63	65	74	548	526	320	292	277	398	343	162	96	17	13
1946	3,466	12	9	24	17	10	19	41	32	44	37	356	290	346	376	306	516	475	337	182	34	3
1947	3,591	15	4	21	19	19	17	17	30	30	28	255	303	371	391	324	573	527	376	213	52	6
1948	3,274	13	11	10	7	19	15	31	30	21	22	185	198	318	358	319	532	517	388	219	59	2
1949	3,402	19	21	34	21	17	9	28	24	20	27	169	215	286	318	300	663	533	426	199	69	4
1950	3,653	14	17	19	26	21	13	28	19	15	28	122	183	262	320	333	620	591	524	361	122	15
1951	3,105	10	13	25	21	19	20	23	23	18	11	91	138	226	337	318	566	474	449	235	78	10
1952	2,326	17	19	25	10	26	16	19	14	15	14	67	149	176	220	227	413	336	311	173	68	11

TABLE 24C.—REPORTED CASES OF TUBERCULOSIS, ALL FORMS, BY YEAR OF REPORT, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1933-1952 (Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1933	2,524	76.1	1,158	70.2	1,366	81.9	1,437	61.0	663	56.1	774	65.8	1,087	113.1	495	105.7	592	120.2	----------
1934	2,047	60.9	904	54.1	1,143	67.7	1,260	52.7	559	46.6	701	58.8	783	80.9	344	72.9	439	88.6	4
1935	1,996	59.0	881	52.4	1,115	65.6	1,133	47.0	508	42.1	625	52.0	863	88.9	373	78.8	490	98.6	----------
1936	3,008	88.5	1,462	86.5	1,546	90.4	1,558	64.2	760	62.5	798	65.8	1,393	143.1	672	141.4	721	144.8	57
1937	2,632	76.7	1,140	66.8	1,492	86.4	1,204	49.1	548	44.6	656	53.5	1,416	144.7	587	122.8	829	165.8	12
1938	2,607	75.0	1,146	66.3	1,461	83.6	1,262	50.7	546	43.8	716	57.5	1,329	135.0	594	123.3	735	146.1	16
1939	2,366	67.0	1,065	60.7	1,262	71.0	1,140	44.9	560	44.2	580	45.7	1,187	119.3	505	103.6	682	134.3	39
1940	2,084	58.3	1,021	57.6	1,063	59.1	930	36.2	493	38.5	437	34.0	1,154	114.9	528	107.5	626	122.1	----------
1941	2,425	66.1	1,209	66.4	1,202	65.0	1,144	43.4	606	46.1	538	40.7	1,267	122.4	603	119.1	664	125.6	14
1942	1,902	51.9	1,040	57.7	826	44.4	932	35.4	580	44.6	352	26.5	934	90.4	460	91.6	474	89.2	36
1943	1,633	44.3	883	49.3	746	39.4	809	30.7	484	37.8	324	23.9	820	78.4	398	78.3	422	78.5	4
1944	1,804	50.2	1,058	62.1	736	38.9	897	34.8	567	46.8	330	24.2	897	87.9	491	99.7	406	76.8	10
1945	3,392	96.5	1,741	106.3	1,651	88.0	1,977	78.7	1,065	92.1	912	67.2	1,415	141.3	676	140.4	739	142.1	----------
1946	3,466	95.2	1,821	102.3	1,645	88.4	2,188	83.9	1,200	93.9	988	74.4	1,278	123.6	621	123.6	657	123.6	----------
1947	3,591	96.6	1,943	106.4	1,648	87.1	2,140	80.4	1,181	90.0	959	71.1	1,451	137.4	762	148.5	689	126.9	----------
1948	3,274	86.2	1,803	96.7	1,470	76.0	1,932	71.0	1,114	83.1	818	59.3	1,342	124.4	689	131.5	652	117.6	----------
1949	3,402	88.0	1,816	95.7	1,586	80.6	2,076	75.0	1,147	84.1	929	66.2	1,325	120.7	668	125.3	657	116.5	1
1950	3,653	89.9	1,919	95.1	1,734	84.8	2,072	69.5	1,172	78.7	900	60.3	1,581	146.6	747	141.7	834	151.2	----------
1951	3,105	75.3	1,854	90.5	1,251	60.3	1,870	61.6	1,141	75.3	729	48.0	1,235	113.5	713	134.1	522	93.8	----------
1952	2,326	55.7	1,385	66.8	941	44.8	1,346	43.7	815	53.0	531	34.5	979	89.4	569	106.3	410	73.2	----------

TABLE 25A.—REPORTED CASES OF TULAREMIA BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept	Oct.	Nov.	Dec.	Not stated
North Carolina	17	13	4		19	10	9		2	1			4	1	3	1	1		4	2	
Alamance	1		1																		
Anson					1		1												1		
Cabarrus	1	1																			
Caswell					3		3						3								
Chowan	1	1																			
Craven	1	1																			
Cumberland	1		1																		
Duplin					1	1							1								
Edgecombe					1	1										1					
Forsyth	1	1			1	1														1	
Granville					2		2								2						
Guilford	1	1																			
Halifax					1		1		1												
Iredell					1	1			1												
Johnston					1	1														1	

Mecklenburg					2	1	1								1				1		
New Hanover	1	1			1	1													1		
Randolph	1	1																			
Rockingham					1	1											1				
Sampson	1		1																		
Stanly	1	1																			
Stokes	1	1																			
Union	2	2																			
Wake	2	2			3	2	1			1				1					1		
Wayne	1		1																		

TABLE 25B.—REPORTED CASES OF TULAREMIA BY YEAR* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	7													1			2	3	1			
1931	3													1	1		1					
1932	10										1			2		1	2	4				
1933	28											1	1	3	1	5	10	5	1			1
1934	14											2		1			8	3				
1935	15											3	2	2	2	2	1	2	1			
1936	10											1		1	2	1	3	2				
1937	13									1		1	2	2	3		1	3				
1938	26											3	3	1	4	3	5	2	2	2		1
1939	26											3	2	3	5	3	7	2	1			
1940	12								1			3	1			1	3	1	2			
1941	22											6	1	1	4	1	1	6	2			
1942	13								1			1		1	2		7		1			
1943	14				1							1	1	1	2	1	2	3	1			1
1944	10													1	3		3	2			1	
1945	27											1	1	2	2	8	6	4	3			
1946	45										1	5	1	4	5	4	8	7	2	4		4
1947	74				1	1		1	2		2	8	4	7	7	8	14	7	8	2		2
1948	42							1		1		4	2	4	4	3	9	8	1	3		2
1949	38					1		1		1		3	3	2	6	1	9	8	3			
1950	25									1		3	1	2	1	4	6	1	5			1
1951	17					1				2		1	2	1	2		2	2	2			2
1952	19			1				1	1			1	3	2	2	1	2	3	1	1		

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 25C.—REPORTED CASES OF TULAREMIA BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	7	0.2	6	0.4	1	0.1	6	0.3	5	0.4	1	0.1	1	0.1	1	0.2	------	------	------
1931	3	0.1	3	0.2	------	------	2	0.1	2	0.2	------	------	1	0.1	1	0.2	------	------	------
1932	10	0.3	5	0.3	5	0.3	9	0.4	4	0.3	5	0.4	1	0.1	1	0.2	------	------	------
1933	28	0.8	20	1.2	8	0.5	23	1.0	16	1.4	7	0.6	4	0.4	3	0.6	1	0.2	1
1934	14	0.4	6	0.4	8	0.5	12	0.5	5	0.4	7	0.6	2	0.2	1	0.2	1	0.2	------
1935	15	0.4	9	0.5	6	0.4	12	0.5	8	0.7	4	0.3	3	0.3	1	0.2	2	0.4	------
1936	10	0.3	4	0.2	6	0.4	8	0.3	3	0.2	5	0.4	2	0.2	1	0.2	1	0.2	------
1937	13	0.4	9	0.5	4	0.2	10	0.4	7	0.6	3	0.2	3	0.3	2	0.4	1	0.2	------
1938	26	0.7	19	1.1	7	0.4	24	1.0	17	1.4	7	0.6	2	0.2	2	0.4	------	------	------
1939	26	0.7	17	1.0	9	0.5	18	0.7	12	0.9	6	0.5	8	0.8	5	1.0	3	0.6	------
1940	12	0.3	8	0.5	4	0.2	11	0.4	7	0.5	4	0.3	1	0.1	1	0.2	------	------	------
1941	22	0.6	13	0.7	9	0.5	18	0.7	11	0.8	7	0.5	4	0.4	2	0.4	2	0.4	------
1942	13	0.4	5	0.3	7	0.4	5	0.2	3	0.2	2	0.2	7	0.7	2	0.4	5	0.9	1
1943	14	0.4	7	0.4	7	0.4	6	0.2	4	0.3	2	0.1	8	0.8	3	0.6	5	0.9	------
1944	10	0.3	5	0.3	5	0.3	7	0.3	5	0.4	2	0.1	3	0.3	------	------	3	0.6	------
1945	27	0.8	15	0.9	12	0.6	22	0.9	13	1.1	9	0.7	5	0.5	2	0.4	3	0.6	------
1946	45	1.2	29	1.6	16	0.9	33	1.3	19	1.5	14	1.1	12	1.2	10	2.0	2	0.4	------
1947	74	2.0	49	2.7	25	1.3	51	1.9	33	2.5	18	1.3	23	2.2	16	3.1	7	1.3	------
1948	42	1.1	22	1.2	20	1.0	32	1.2	16	1.2	16	1.2	10	0.9	6	1.1	4	0.7	------
1949	38	1.0	24	1.3	14	0.7	31	1.1	20	1.5	11	0.8	7	0.6	4	0.8	3	0.5	------
1950	25	0.6	13	0.6	12	0.6	19	0.6	11	0.7	8	0.5	6	0.6	2	0.4	4	0.7	------
1951	17	0.4	10	0.5	7	0.3	13	0.4	8	0.5	5	0.3	4	0.4	2	0.4	2	0.4	------
1952	19	0.5	10	0.5	9	0.4	10	0.3	7	0.5	3	0.2	9	0.8	3	0.6	6	1.1	------

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 26A.—REPORTED CASES OF TYPHOID FEVER BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	44	31	13		41	25	16				2	1	7	6	6	8	5	2	3	1	
Alamance					2	2										2					
Alexander	1	1																			
Beaufort	2	1	1		2		2								1		1				
Brunswick					1	1											1				
Buncombe	4	3	1		1		1						1								
Burke	1	1																			
Caldwell	1	1			1	1								1							
Carteret	1	1			1	1													1		
Cherokee					1	1											1				
Columbus					1	1								1							
Cumberland	1	1																			
Davidson	1	1																			
Davie					1	1											1				
Durham					1	1											1				
Edgecombe					1		1						1								

Forsyth	3	3																		
Franklin					2	2					1					1				
Gaston	2	2			1	1									1					
Graham					1	1													1	
Granville					1	1										1				
Greene					1		1								1					
Guilford	4	3	1																	
Halifax					4		4								2			2		
Henderson	1	1																		
Hoke	1		1																	
Iredell	2	2			1	1										1				
Madison	2	2			1	1							1							
Mecklenburg	1	1			1	1										1				
Nash	1	1			2		2									1			1	
New Hanover	2	1	1																	
Orange	1	1																		
Person					1	1										1				
Pitt	7		7																	
Polk					1	1								1						
Richmond	1		1																	
Robeson					7	3	4				1	1	4	1						
Rockingham					1	1								1						
Rowan	2	2																		
Sampson	1	1																		
Wake					1	1													1	
Wayne	1	1																		
Wilson					2	1	1							1	1					

TABLE 26B.—REPORTED CASES OF TYPHOID FEVER BY YEAR* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	1,000	1	5	7	24	20	33	31	41	45	39	174	160	116	81	64	78	38	16	8	4	15
1931	991	1	3	8	20	31	32	29	44	38	37	165	170	103	78	53	77	56	29	7	2	8
1932	823	------	4	8	15	28	29	36	28	39	27	136	139	92	60	43	55	48	20	7	2	7
1933	684	1	6	9	16	20	24	24	20	26	32	109	112	85	52	38	52	27	12	8	3	8
1934	464	3	1	1	5	7	20	18	20	18	17	90	67	53	43	27	42	15	8	------	------	9
1935	645	1	------	7	12	21	16	19	27	22	21	107	91	97	50	33	53	31	12	6	2	17
1936	518	2	4	9	5	15	20	15	20	18	15	79	59	59	52	46	38	29	13	7	------	13
1937	493	4	6	5	16	18	22	16	22	21	15	65	62	55	44	41	43	18	12	2	------	6
1938	485	1	4	6	15	15	16	18	21	16	19	67	72	51	37	29	35	28	13	5	------	17
1939	343	2	6	10	9	18	16	13	9	8	13	53	48	32	33	22	28	9	7	1	1	5
1940	238	1	1	5	8	9	6	6	4	7	6	32	22	33	35	19	16	13	5	6	------	4
1941	197	2	1	4	11	7	15	6	10	2	4	26	27	19	19	12	13	8	3	2	------	6
1942	181	1	3	6	12	7	6	10	7	5	10	30	21	12	10	7	16	11	4	1	------	2
1943	111	1	1	1	4	3	7	10	5	4	4	11	15	9	8	2	13	6	2	1	1	3
1944	122	1	------	1	2	6	5	5	4	7	6	30	13	9	5	3	14	5	2	1	1	2
1945	82	1	------	4	5	1	2	------	4	1	2	14	13	6	4	8	11	4	2	------	------	--------
1946	54	------	1	------	------	3	1	3	1	6	1	8	5	2	6	1	7	3	4	2	------	--------
1947	47	1	------	2	3	4	4	1	1	1	3	8	2	1	3	4	4	5	------	------	------	--------
1948	59	1	6	1	2	3	5	4	3	3	------	6	8	5	3	3	4	1	------	1	------	--------
1949	53	------	3	2	2	2	1	1	1	3	1	6	7	2	4	5	4	6	2	1	------	--------
1950	58	------	3	1	2	2	2	4	------	2	------	9	7	6	6	5	3	4	1	------	1	--------
1951	44	------	1	1	4	4	3	------	3	1	1	2	4	4	4	4	3	3	1	1	------	--------
1952	41	------	------	2	------	3	2	5	------	------	------	5	4	------	5	4	6	------	1	3	1	--------

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 26C.—REPORTED CASES OF TYPHOID FEVER BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952 (Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	1,000	31.5	545	34.6	454	28.5	566	25.3	306	27.3	260	23.3	431	46.1	237	52.1	194	40.4	3
1931	991	30.9	532	33.4	459	28.5	609	26.9	322	28.3	287	25.4	382	40.6	210	45.9	172	35.7	----------
1932	823	25.2	444	27.4	379	23.1	517	22.3	289	24.9	228	19.8	306	32.2	155	33.5	151	31.6	----------
1933	684	20.6	385	23.3	299	17.9	365	15.5	213	18.0	152	12.9	316	32.9	170	36.3	146	29.6	3
1934	464	13.8	270	16.2	194	11.5	275	11.5	172	14.3	103	8.6	189	19.5	98	20.8	91	18.4	----------
1935	645	19.1	382	22.7	263	15.5	346	14.4	213	17.6	133	11.1	298	30.7	169	35.7	129	26.0	1
1936	518	15.2	292	17.3	226	13.2	306	12.6	167	13.7	139	11.5	209	21.5	123	25.9	86	17.3	3
1937	493	14.4	264	15.5	229	13.3	299	12.2	164	13.4	135	11.0	191	19.5	98	20.5	93	18.6	3
1938	485	14.0	279	16.2	206	11.8	299	12.0	165	13.2	134	10.8	182	18.5	112	23.2	70	13.9	4
1939	343	9.7	189	10.8	148	8.3	221	8.7	123	9.7	98	7.7	116	11.7	66	13.5	50	9.8	6
1940	238	6.7	137	7.7	101	5.6	161	6.3	91	7.1	70	5.4	77	7.7	46	9.4	31	6.0	----------
1941	197	5.4	112	6.2	84	4.5	106	4.0	58	4.4	48	3.6	90	8.7	54	10.7	36	6.8	1
1942	181	4.9	103	5.7	77	4.1	113	4.3	62	4.8	51	3.8	67	6.5	41	8.2	26	4.9	1
1943	111	3.0	64	3.6	45	2.4	66	2.5	35	2.7	31	2.3	43	4.1	29	5.7	14	2.6	2
1944	122	3.4	77	4.5	45	2.4	88	3.4	57	4.7	31	2.3	34	3.3	20	4.1	14	2.6	----------
1945	82	2.3	45	2.7	37	2.0	52	2.1	28	2.4	24	1.8	30	3.0	17	3.5	13	2.5	----------
1946	54	1.5	35	2.0	18	1.0	32	1.2	23	1.8	9	0.7	21	2.0	12	2.4	9	1.7	1
1947	47	1.3	28	1.5	19	1.0	26	1.0	12	0.9	14	1.0	21	2.0	16	3.1	5	0.9	----------
1948	59	1.6	34	1.8	25	1.3	43	1.6	21	1.6	22	1.6	16	1.5	13	2.5	3	0.5	----------
1949	53	1.4	29	1.5	24	1.2	28	1.0	15	1.1	13	0.9	24	2.2	14	2.6	10	1.8	1
1950	58	1.4	33	1.6	25	1.2	45	1.5	23	1.5	22	1.5	13	1.2	10	1.9	3	0.5	----------
1951	44	1.1	23	1.1	21	1.0	31	1.0	18	1.2	13	0.9	13	1.2	5	0.9	8	1.4	----------
1952	41	1.0	20	1.0	21	1.0	25	0.8	11	0.7	14	0.9	16	1.5	9	1.7	7	1.2	----------

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 27A.—REPORTED CASES OF TYPHUS FEVER, MURINE, BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	19	16	3		12	11	1		1					1	2	4	1	3			
Anson	1	1																			
Caldwell					1	1									1						
Caswell	1		1																		
Catawba					1	1			1												
Chatham	1	1																			
Clay	1	1																			
Duplin	1	1																			
Guilford					3	3									1	1		1			
Iredell	1	1																			
Johnston	2	2																			
Lee	3	3																			
Mecklenburg	3	3			2	1	1									1		1			
Nash					1	1												1			
New Hanover					1	1											1				
Randolph					1	1										1					
Richmond	2	1	1																		
Rowan	1	1																			
Vance	1	1			1	1										1					
Wayne					1	1								1							
Wilson	1		1																		

Table 27B.—Reported Cases of Typhus Fever, Murine, by Year* and Age, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	16						1			2		3		2	3		1	3	1			
1931	14					1						2	1		1	2	1	2	3			1
1932	24					1			1	1		2	3	1	4		6	1	2			2
1933	46					2	1			1	1	2	5	6	6	3	9	5	3	1		1
1934	23											2	3	9		1	5	1	1		1	
1935	51						1			1		3	3	9	6	3	10	9	3			3
1936	33								1		1	1	6	1	4	3	8	4	2	2		
1937	68					1			2	2	1	2	6	5	7	11	9	15	3	4		
1938	81			2					1	3		5	8	14	8	8	13	8	6	4		1
1939	122			2	1	4	1	2	2	1	1	7	12	18	16	10	17	15	7	2	1	3
1940	65					1	1	1	1		1	4	7	10	4	9	10	6	3	4		3
1941	78				1					1		2	14	11	12	7	8	7	5	2		8
1942	100				1			1	2			11	8	9	9	12	21	10	8	4		4
1943	157	1			1		1				1	8	19	11	20	14	27	27	12	4		11
1944	236				1	4		1	1	3	2	8	22	18	22	24	46	40	24	9		11
1945	159			1		1	1	1	2	3		13	9	8	21	16	32	18	20	5		8
1946	82				1	2		1	1	1	2	9	9	7	5	7	13	9	8	4	1	2
1947	51			1		1	2	1	1	1	1	3	2		4	5	10	6	5	5	2	1
1948	45					1		3	2	1	1	3	2	2	2	4	8	5	7	3		1
1949	26				1				1	1		4	2	2	1	5	8				1	
1950	12							1				1		1	3		3	2	1			
1951	19		1	1				1	1	2		3	2		2	3	2					1
1952	12			1								3	1	2		1	2		2			

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 27C.—REPORTED CASES OF TYPHUS FEVER, MURINE, BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	16	0.5	10	0.6	6	0.4	15	0.7	10	0.9	5	0.4	1	0.1	------	------	1	0.2	------
1931	14	0.4	13	0.8	1	0.1	14	0.6	13	1.1	1	0.1	------	------	------	------	------	------	------
1932	24	0.7	12	0.7	12	0.7	21	0.9	9	0.8	12	1.0	3	0.3	3	0.6	------	------	------
1933	46	1.4	31	1.9	15	0.9	42	1.8	29	2.5	13	1.1	4	0.4	2	0.4	2	0.4	------
1934	23	0.7	10	0.6	13	0.8	18	0.8	7	0.6	11	0.9	5	0.5	3	0.6	2	0.4	------
1935	51	1.5	32	1.9	19	1.1	48	2.0	30	2.5	18	1.5	3	0.3	2	0.4	1	0.2	------
1936	33	1.0	27	1.6	6	0.4	33	1.4	27	2.2	6	0.5	------	------	------	------	------	------	------
1937	68	2.0	53	3.1	15	0.9	57	2.3	46	3.7	11	0.9	10	1.0	7	1.5	3	0.6	1
1938	81	2.3	58	3.4	22	1.3	71	2.9	53	4.3	18	1.4	9	0.9	5	1.0	4	0.8	1
1939	122	3.5	75	4.3	41	2.3	104	4.1	67	5.3	37	2.9	12	1.2	8	1.6	4	0.8	6
1940	65	1.8	48	2.7	17	0.9	54	2.1	38	3.0	16	1.2	11	1.1	10	2.0	1	0.2	------
1941	78	2.1	56	3.1	20	1.1	66	2.5	48	3.7	18	1.4	10	1.0	8	1.6	2	0.4	2
1942	100	2.7	63	3.5	32	1.7	86	3.3	58	4.5	28	2.1	9	0.9	5	1.0	4	0.8	5
1943	157	4.3	92	5.1	64	3.4	144	5.5	86	6.7	58	4.3	12	1.1	6	1.2	6	1.1	1
1944	236	6.6	158	9.3	73	3.9	217	8.4	149	12.3	68	5.0	14	1.4	9	1.8	5	0.9	5
1945	159	4.5	90	5.5	67	3.6	141	5.6	79	6.8	62	4.6	16	1.6	11	2.3	5	1.0	2
1946	82	2.3	43	2.4	38	2.0	72	2.8	38	3.0	34	2.6	9	0.9	5	1.0	4	0.8	1
1947	51	1.4	29	1.6	22	1.2	46	1.7	26	2.0	20	1.5	5	0.5	3	0.6	2	0.4	------
1948	45	1.2	23	1.2	22	1.1	37	1.4	18	1.3	19	1.4	8	0.7	5	1.0	3	0.5	------
1949	26	0.7	20	1.1	6	0.3	22	0.8	17	1.2	5	0.4	4	0.4	3	0.6	1	0.2	------
1950	12	0.3	6	0.3	6	0.3	10	0.3	6	0.4	4	0.3	2	0.2	------	------	2	0.4	------
1951	19	0.5	5	0.2	14	0.7	16	0.5	3	0.2	13	0.9	3	0.3	2	0.4	1	0.2	------
1952	12	0.3	7	0.3	5	0.2	11	0.4	7	0.5	4	0.3	1	0.1	------	------	1	0.2	------

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 28A.—REPORTED CASES OF UNDULANT FEVER BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	28	24	4		21	15	6		2	2	1		2	1	1	1	1	6	1	1	2
Bertie					1	1															1
Chatham					1	1															1
Chowan					3		3						1					2			
Cleveland	1	1																			
Craven	3	3																			
Currituck	1	1																			
Gaston	1	1			3	3			1	2											
Gates	2	1	1																		
Guilford	3	3																			
Halifax	1	1			2	1	1								1					1	
Iredell	2	2			2	2			1								1				
Lenoir	1	1																			
Lincoln	1	1																			
Martin	2	2			1	1								1							
Mecklenburg	2	2			1	1										1					

TABLE 28A.—UNDULANT FEVER—*Continued*

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Nash	2	2			1	1												1			
New Hanover	1		1																		
Onslow	1		1																		
Pasquotank					1		1											1			
Perquimans	1	1																			
Pitt					2	1	1				1								1		
Transylvania					2	2												2			
Warren					1	1							1								
Wayne	1		1																		
Wilson	1	1																			
Yancey	1	1																			

TABLE 28B.—REPORTED CASES OF UNDULANT FEVER BY YEAR* AND AGE, 1932-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1932	4											1	1						1			1
1933	15													3	3	2	5		1			1
1934	19											2	1	1	2	3	5	1	2			2
1935	31						1			1		2	1	5	5	3	6	5	1	1		
1936	23											2		2	5	2	4	3	3	1		1
1937	26											1	2	2	4	2	6	2	2	1		4
1938	29											4	2	6	4	3	5	1	3			1
1939	25							1	1		1		2	2	3	4	3	6	1			1
1940	11										1			2	4	1		2				1
1941	12													1	4	1	2	3				1
1942	9											1	1		1	1	1	1				3
1943	7												1	1			3	1		1		
1944	14												1	3	1	1	4	3	1			
1945	15											1	1	1	2	3	3	4				
1946	9													1	1	2	1	3		1		
1947	21					1		1				1		1	2	2	9	2	1			1
1948	15											1	1	2		1	5	3	1	1		
1949	25		1	1									1	3	3	2	7	4		1	1	1
1950	21							1				1			2	2	7	3	4	1		
1951	28								1			1		4		2	13	2	5			
1952	21													2	2	1	8	6	1			1

* 1932-1948 by year of report; 1949-1952 by year of onset.

TABLE 28C.—REPORTED CASES OF UNDULANT FEVER BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1932-1952
(Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1932	4	0.1	2	0.1	2	0.1	4	0.2	2	0.2	2	0.2							
1933	15	0.5	8	0.5	7	0.4	13	0.6	8	0.7	5	0.4	2	0.2			2	0.4	
1934	19	0.6	16	1.0	3	0.2	16	0.7	14	1.2	2	0.2	3	0.3	2	0.4	1	0.2	
1935	31	0.9	20	1.2	11	0.6	22	0.9	13	1.1	9	0.7	9	0.9	7	1.5	2	0.4	
1936	23	0.7	15	0.9	8	0.5	20	0.8	13	1.1	7	0.6	3	0.3	2	0.4	1	0.2	
1937	26	0.8	15	0.9	11	0.6	25	1.0	15	1.2	10	0.8	1	0.1			1	0.2	
1938	29	0.8	18	1.0	11	0.6	20	0.8	14	1.1	6	0.5	9	0.9	4	0.8	5	1.0	
1939	25	0.7	13	0.7	10	0.6	19	0.7	9	0.7	10	0.8	4	0.4	4	0.8			2
1940	11	0.3	8	0.5	3	0.2	9	0.4	6	0.5	3	0.2	2	0.2	2	0.4			
1941	12	0.3	9	0.5	3	0.2	9	0.3	7	0.5	2	0.2	3	0.3	2	0.4	1	0.2	
1942	9	0.2	8	0.4	1	0.1	7	0.3	6	0.5	1	0.1	2	0.2	2	0.4			
1943	7	0.2	5	0.3	2	0.1	7	0.3	5	0.4	2	0.1							
1944	14	0.4	10	0.6	4	0.2	12	0.5	8	0.7	4	0.3	2	0.2	2	0.4			
1945	15	0.4	12	0.7	3	0.2	13	0.5	11	1.0	2	0.1	2	0.2	1	0.2	1	0.2	
1946	9	0.2	5	0.3	4	0.2	8	0.3	5	0.4	3	0.2	1	0.1			1	0.2	
1947	21	0.6	12	0.7	9	0.5	17	0.6	10	0.8	7	0.5	4	0.4	2	0.4	2	0.4	
1948	15	0.4	10	0.5	5	0.3	11	0.4	6	0.4	5	0.4	4	0.4	4	0.8			
1949	25	0.6	13	0.7	12	0.6	21	0.8	10	0.7	11	0.8	4	0.4	3	0.6	1	0.2	
1950	21	0.5	17	0.8	4	0.2	17	0.6	13	0.9	4	0.3	4	0.4	4	0.8			
1951	28	0.7	22	1.1	6	0.3	24	0.8	19	1.3	5	0.3	4	0.4	3	0.6	1	0.2	
1952	21	0.5	13	0.6	8	0.4	15	0.5	9	0.6	6	0.4	6	0.5	4	0.7	2	0.4	

* 1932-1948 by year of report; 1949-1952 by year of onset.

TABLE 29A.—REPORTED CASES OF WHOOPING COUGH BY COUNTY AND COLOR, 1951 AND 1952, WITH MONTH OF ONSET, 1952
(Place of Residence)

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
North Carolina	1,961	1,207	739	15	471	375	92	4	56	46	59	60	64	26	22	29	43	13	28	7	18
Alamance	2	2			9	8	1		1		2	4		2							
Alexander	3	3																			
Alleghany					1	1					1										
Anson	5	5																			
Ashe	1	1			13	13				2	11										
Avery	4	4																			
Beaufort	7	1	6																		
Bertie	3	1	2																		
Bladen	2	1	1																		
Brunswick	23	18	4	1	1	1													1		
Buncombe	54	53	1		59	59			4	9	13	13		5	3		11	1			
Burke	11	10	1		2	2			1				1								
Cabarrus	35	26	9		1	1													1		
Caldwell	4	3	1		1	1			1												
Camden	1	1			2	2			1												1

TABLE 29A.—WHOOPING COUGH—*Continued*

Area	1951 Onsets				1952 Onsets																
	Color				Color				Month of onset												
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Carteret	4	4																			
Caswell	18		18																		
Catawba	2	2																			
Chatham	12	3	9																		
Cherokee	4	4																			
Chowan	20	18	2																		
Clay	1	1																			
Cleveland	41	30	11		9	5	4		1		1	1	1	1				4			
Columbus	7	4	3		2	2										2					
Craven	1	1			5	5									1	4					
Cumberland	6		6																		
Currituck	4	4			1		1					1									
Dare	1	1			1	1				1											
Davidson	6	6																			
Davie	6	6																			
Duplin	1		1		5	4	1			1			3	1							
Durham	211	113	98		21	19	2		3		4	1	1	1		2	1	1			7
Edgecombe	7	6	1		5	5						1			3	1					
Forsyth	64	54	10		5	2	3						1			1			3		
Franklin	16	13	3		12	5	7				2	1	3	4		2					

Gaston	38	37	1		19	19							4	1			1		8	5	
Gates	23	8	15																		
Graham	41	41			37	36		1	21	10			3	3							
Granville	35	10	25																		
Guilford	22	17	5		24	9	15			1	1	1	2			12	4	2			1
Halifax	20	9	11		19	4	15		6	2	4	4	2		1						
Harnett	14	13	1		1		1		1												
Haywood	1	1			20	20										2	18				
Henderson	8	8			1	1			1												
Hertford	6	2	4																		
Hyde	1		1																		
Iredell	25	17	7	1	2	2											2				
Jackson	3	3			1		1												1		
Johnston	7	7																			
Lee	1	1																			
Lenoir	2	2																			
McDowell	4	4																			
Macon	1	1			4	4						3	1								
Madison	32	32			2	2			2												
Martin	4	1	3																		
Mecklenburg	134	83	45	6	35	24	11			2	4	6	4		3	2	1	3	10		
Mitchell	16	15		1	1			1				1									
Montgomery	5		5																		
Moore	13	11	2		2	1	1		1	1											
Nash	28	25	3		4	1	3			2		1	1								
New Hanover	61	42	18	1	11	7	4		1			2	3	2	3						
Onslow	10	10			2	2											2				
Orange	3	3			22	22			2	6	1	11	2								
Pamlico	3	3			13	13			1		2	3	7								
Pasquotank	19	5	14		10	4	6		7	2			1								

TABLE 29A.—WHOOPING COUGH—*Continued*

Area	1951 Onsets				1952 Onsets															
	Color				Color				Month of onset											
	Total	White	Other	Not stated	Total	White	Other	Not stated	Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Not stated
Person	8	7	1		1	1					1										
Randolph	12	8	4																		
Richmond	11	10		1																	
Robeson	46	2	43	1	10	3	6	1				2			1		1	1	4	1	
Rockingham	5	5			3	1	2						1				2				
Rowan	35	24	11		1		1											1			
Rutherford	11	7	4																		
Sampson	7	5	2																		
Scotland	26	12	14		1	1					1										
Stanly	18	18			1	1														1	
Surry	8	8			2	2			1		1										
Swain	2		2		7	6	1			1	5										1
Tyrrell	18	9	9																		
Union	1	1																			
Vance	4	3	1																		
Wake	477	239	238		14	9	5			4	5	3		2							
Warren	7		7																		
Washington	66	16	48	2																	
Watauga	1	1			45	44		1		2		1	22	4	7	1					8
Wilson	22	13	8	1	1		1						1								
Yadkin	3	3																			
Yancey	6	6																			

TABLE 29B.—REPORTED CASES OF WHOOPING COUGH BY YEAR* AND AGE, 1930-1952 (Place of Residence)

Year	All ages	Under 1 Yr.	1 Yr.	2 Yrs.	3 Yrs.	4 Yrs.	5 Yrs.	6 Yrs.	7 Yrs.	8 Yrs.	9 Yrs.	10-14 Yrs.	15-19 Yrs.	20-24 Yrs.	25-29 Yrs.	30-34 Yrs.	35-44 Yrs.	45-54 Yrs.	55-64 Yrs.	65-74 Yrs.	75 and over	Not stated
1930	10,093	810	713	913	986	941	797	1,105	691	529	384	596	69	25	10	9	3	2	1	------	1	1,508
1931	6,992	575	567	706	773	797	689	857	677	412	249	372	51	16	5	1	5	2	1	------	------	237
1932	12,137	1,029	968	1,330	1,360	1,387	1,209	1,417	1,150	843	460	731	99	19	14	10	7	------	------	------	------	104
1933	7,155	585	548	698	750	711	661	945	643	499	326	609	70	14	8	5	5	2	2	1	------	73
1934	14,299	1,211	1,133	1,403	1,512	1,597	1,331	1,717	1,350	973	582	1,091	128	24	12	6	6	1	------	------	------	222
1935	10,075	936	775	997	1,089	1,129	911	1,225	993	660	403	697	74	32	6	9	5	2	1	1	------	130
1936	1,566	169	126	164	168	175	142	202	137	106	51	102	8	1	------	2	4	1	------	------	------	8
1937	7,782	717	710	803	843	848	748	915	713	534	292	430	49	15	12	10	4	2	1	------	------	136
1938	14,908	1,405	1,210	1,466	1,671	1,533	1,337	1,690	1,391	1,110	672	999	130	24	20	13	7	4	3	2	------	221
1939	9,321	876	830	1,035	1,120	1,091	895	1,063	755	541	364	522	56	9	9	9	2	2	2	------	------	140
1940	5,500	528	442	538	588	599	567	691	532	382	219	306	15	6	1	5	6	------	------	------	------	75
1941	10,932	1,031	829	1,264	1,318	1,171	1,061	1,338	1,003	734	366	588	52	16	11	9	7	6	1	4	------	123
1942	5,557	569	327	663	675	603	524	714	543	318	186	322	38	5	7	5	5	------	------	------	------	53
1943	8,408	863	496	1,001	1,036	981	824	955	806	583	289	398	37	14	8	4	5	1	------	------	1	106
1944	6,355	691	380	876	798	729	645	641	521	376	208	321	41	8	7	1	3	1	------	------	------	108
1945	5,834	613	348	774	682	715	554	663	511	359	230	315	24	10	6	6	4	3	------	------	------	17
1946	3,391	376	202	283	419	409	303	362	338	215	142	210	17	3	6	2	------	------	------	------	------	4
1947	2,983	439	172	341	384	334	290	270	243	185	115	171	10	2	3	3	2	------	1	2	------	16
1948	1,976	332	193	214	222	179	195	164	130	119	76	115	17	2	2	2	2	------	------	------	------	12
1949	1,398	195	138	153	158	139	115	143	93	82	52	109	11	2	2	------	1	------	------	------	------	5
1950	3,352	457	297	375	392	345	276	280	242	202	146	265	31	2	4	2	2	1	1	------	------	32
1951	1,961	294	193	220	226	217	157	163	98	118	76	161	15	4	1	1	2	------	------	------	------	15
1952	471	93	59	47	50	37	35	32	34	25	18	31	2	------	2	------	------	------	------	------	------	6

* 1930-1948 by year of report; 1949-1952 by year of onset.

TABLE 29C.—REPORTED CASES OF WHOOPING COUGH BY YEAR*, COLOR, AND SEX WITH RATES PER 100,000 POPULATION, 1930-1952 (Place of Residence)

Year	All classes						White						Other						Not stated sex and/or color
	Total		Male		Female		Total		Male		Female		Total		Male		Female		
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate	
1930	10,093	318.3	4,249	269.6	4,436	278.1	7,561	338.1	3,636	324.4	3,776	338.6	1,194	127.7	574	126.2	620	129.2	1,408
1931	6,992	218.0	3,306	207.4	3,520	281.2	5,975	263.6	2,871	252.7	3,053	270.0	810	86.2	384	83.9	426	88.3	207
1932	12,137	371.7	5,739	353.6	6,151	374.6	10,424	450.5	5,037	434.3	5,387	466.8	1,220	128.3	567	122.4	653	133.9	493
1933	7,155	215.6	3,531	214.0	3,606	216.1	5,649	239.6	2,770	234.4	2,864	243.6	1,111	115.6	548	117.0	563	114.3	395
1934	14,299	425.7	6,873	411.4	7,313	433.1	11,804	493.5	5,646	471.0	6,055	507.5	2,274	235.1	1,110	235.3	1,154	232.9	221
1935	10,075	298.1	4,985	296.5	5,090	299.6	8,637	358.4	4,304	356.4	4,333	360.4	1,366	140.8	646	136.4	720	144.9	72
1936	1,566	46.1	724	42.8	842	49.2	1,240	51.1	573	47.2	667	55.0	323	33.2	150	31.6	173	34.7	3
1937	7,782	226.7	3,710	217.5	4,072	235.9	6,972	284.1	3,322	270.6	3,650	297.6	744	76.1	353	73.8	391	78.2	66
1938	14,908	428.9	7,087	410.3	7,821	447.4	11,876	476.8	5,685	456.4	6,191	497.2	2,946	299.2	1,364	283.1	1,582	314.5	86
1939	9,321	263.9	4,325	246.4	4,688	263.9	6,888	271.5	3,318	261.6	3,570	281.4	2,125	213.6	1,007	206.6	1,118	220.2	308
1940	5,500	154.0	2,676	150.9	2,822	156.9	4,493	175.0	2,191	170.9	2,302	179.0	1,006	100.2	485	98.7	521	101.6	2
1941	10,932	297.9	5,061	278.0	5,633	304.5	8,666	328.9	4,117	313.4	4,549	344.3	2,028	196.0	944	186.4	1,084	205.1	238
1942	5,557	151.6	2,580	143.1	2,819	151.4	4,222	160.4	2,017	155.0	2,205	165.7	1,177	113.9	563	112.1	614	115.5	158
1943	8,408	228.3	3,671	205.1	4,714	249.0	7,132	270.5	3,159	246.5	3,973	293.1	1,253	119.8	512	100.7	741	137.8	23
1944	6,355	176.8	2,961	173.8	3,193	168.9	4,425	171.9	2,142	176.8	2,283	167.6	1,729	169.4	819	166.3	910	172.2	201
1945	5,834	166.0	2,731	166.8	3,049	162.5	4,427	176.2	2,085	180.3	2,342	172.6	1,353	135.1	646	134.2	707	135.9	54
1946	3,391	93.1	1,634	91.8	1,740	93.5	2,694	103.3	1,312	102.7	1,382	104.0	680	65.8	322	64.1	358	67.4	17
1947	2,983	80.2	1,421	77.8	1,562	82.5	1,948	73.2	935	71.2	1,013	75.1	1,035	98.0	486	94.7	549	101.1	----------
1948	1,976	52.0	953	51.1	1,023	52.9	1,366	50.2	659	49.2	707	51.3	594	55.1	284	54.2	310	55.9	16
1949	1,398	36.2	658	34.7	740	37.6	1,071	38.7	497	36.4	574	40.9	321	29.3	159	29.8	162	28.7	6
1950	3,352	82.5	1,619	80.3	1,731	84.7	2,519	84.4	1,195	80.2	1,324	88.7	817	75.7	417	79.1	400	72.5	16
1951	1,961	47.6	935	45.7	1,024	49.3	1,207	39.8	589	38.9	617	40.6	739	67.9	338	63.6	401	72.1	15
1952	471	11.3	214	10.3	257	12.2	375	12.2	175	11.4	200	13.0	92	8.4	36	6.7	56	10.0	4

* 1930-1948 by year of report; 1949-1952 by year of onset.

UNIVERSITY OF N.C. AT CHAPEL HILL
00015568921

This **BOOK** may be kept out **TWO WE**
ONLY, and is subject to a fine of
CENTS a day thereafter. It is **DUE** o
DAY indicated below: